



## SEQUENTIAL CHANGES IN COAGULATION FACTORS AND CELLULAR COMPONENTS OF ACD-STORED BLOOD.

Pramoat Teowsiri, B.Sc. (M.T.)\*

Lumduan Changlaw, B.Sc. (M.T.)\*

Suchada Tawarat, B.Sc. (M.T.)\*

Panja Kulapongs, M.D.\*

### ABSTRACT.

Sequential changes in blood cell components and coagulation factors during short-term storage in blood bank condition was studied in 20 blood samples stored at 4°C in ACD solution. The recovery rates of leukocytes, leukocyte viability and platelets were comparable to that observed by others in both ACD and CPD blood. The recovery rate of leukocytes of  $85.45 \pm 18.30\%$  at one week and 80% at 10 days of storage were similar to that of the CPD blood. Most of the leukocytes disintegrated were PMNs. The finding of higher recovery rate of platelets (46% at 2 weeks of storage) in our ACD blood is probably due to the fact that spontaneous clumping of platelets occurred less rapidly in ACD blood. Factor VIII activity dropped rapidly within the first few hours and only  $33.90 \pm 8.68\%$  of its activity was detected during the first day of blood collection. Factor V activity decreased more gradually to only 40% of the original activity by day 7. Factor VII activity increased gradually within the first week of storage due to the cold-promoted enhancement of its activity via plasma kallikrein and contact activation.

---

\*Department of Clinical Microscopy, Faculty of Associated Medical Sciences,  
Chiang Mai University.

## INTRODUCTION.

Sodium citrate was first introduced as a useful anticoagulant for blood collection and transfusion by Lewisohn in 1915<sup>(1)</sup>. The addition of dextrose to this anticoagulant further extended the usable time for the collected blood<sup>(2)</sup>. The replacement of some citrate with citric acid by Loutit et al in 1943<sup>(3)</sup> resulting in a solution almost identical to the present Acid-citrate - Dextrose (ACD) solutions. Although other anticoagulants, such as heparin and chelating agents, have been used for special purposes, ACD has remained the standard blood preservative for many years. The ACD preserved blood has a usable span of only 21 days before decreasing red cell viability made it unsatisfactory for transfusion purposes. In 1957, Gibson et al<sup>(4)</sup> introduced citrate-phosphate - dextrose (CPD) solution which has a higher pH, lower citrate concentration and a more isotonic environment than ACD solutions. Results of several studies indicated that the percentage of red cell surviving at 28 days of storage in CPD was significantly greater than in ACD solutions. In additions, the presence of sodium phosphate in CPD solution results in better conservation of 2.3 DPG than occurs in ACD stored blood and thus maintains better oxygen transport efficiency.

Although CPD is considered to be the best routine blood preservative which should prolong the usable span of stored red cells, a few limitations to its widespread use exist. Prolongation of the storage time by CPD results in higher plasma levels of ammonia, potassium and free hemoglobin (5-7) and greater amount of intracellular potassium liberated from stored red cells that are lysed in the immediate posttransfusion period<sup>(7)</sup>. The concentration of plasma sodium is greater in CPD stored blood due to the use of sodium citrate and sodium phosphate CPD solution<sup>(5)</sup>. These substances could result in potentially deleterious effects in patients with hepatic and renal diseases. In addition, the formation of microaggregates of leukocytes and platelets is found to be greater in CPD than in ACD stored blood<sup>(8)</sup> which is probably due to the higher pH of the former<sup>(9)</sup>.

In practice, blood is stored in ACD solution and the storage time is rarely longer than 2 weeks in our blood bank. We are reporting the sequential changes in coagulation factors, chemical and blood cell components occurred in ACD stored blood in such short-term storage.

## MATERIALS AND METHODS.

Blood samples were collected from 20 healthy male adults according

to the guidelines of the American Association of Blood Bank<sup>(10)</sup>. Approximately 400 ml of blood were collected into 10 plastic bags and 10 glass bottles containing 67 ml and 100 ml of ACD solution respectively. After thorough mixing, whole blood samples were aliquoted and stored at 4°C in sterile plastic or glass containers accordingly. Each blood samples were analyzed immediately following collection and on days 1, 2, 3, 4, 5, 6, 9 and 13. White blood cell count, differential leukocyte count, leukocyte viability test (trypan blue dye exclusion test) and platelet count were determined on all 20 blood samples. Plasma concentrations of sodium and potassium were determined by atomic absorption spectroscopic technique<sup>(11)</sup>. Plasma levels of coagulation factors V, VII, VIII, IX, X and XII were determined<sup>(12,17)</sup> on 10 blood samples collected and stored in plastic containers.

## RESULTS

The baseline values of the blood cell components, coagulation factors and their sequential changes during storage at 4°C were summarized and shown in Table 1 and Figure 1. The recovery rate of leukocytes of 85.45 ± 18.30% at one week was slightly low but the rate of 80% at 10 days of storage was comparable to that of

the CPD blood<sup>(18)</sup>. Most of the leukocytes disintegrated were PMNs. At 2 weeks 66.0 ± 3.13% of the remaining leukocytes were still viable, and the recovery rate of platelets was 46%. The plasma K<sup>+</sup> concentration rise gradually from the baseline value of 3.30 ± 0.8 mEq/L to 15.5 ± 2.95 mEq/L after 2 weeks of storage comparable to that observed by others for both ACD and CPD blood<sup>(7, 19)</sup>. Significantly low levels of factor VIII (33.90 ± 8.68%), IX (36.5 ± 5.08%) and XII (26.4 ± 7.59%) were observed in the first day of blood collection. Factor V activity decreased gradually to 40% of the original activity by day 7. Factor VII activity increased 3 folds during the first week of storage.

## COMMENTS

The plasma K<sup>+</sup> concentration of our ACD blood increased in direct proportion to the storage time (Figure 1). The magnitude of the increment was comparable to that observed by others for both ACD and CPD blood<sup>(5-7, 20-24)</sup>. Normally red cells contain approximately 100 mEq K<sup>+</sup>/L. of packed red cells and plasma contains only 3-5 mEq K<sup>+</sup>/L.. Stored red cells have been known to lose K<sup>+</sup> into the surrounding plasma<sup>(25)</sup> resulting in increased plasma K<sup>+</sup> concentration and decreased K<sup>+</sup> content of red cells.

The increased in plasma  $K^+$ , LDH and free hemoglobin levels reflect either red cell hemolysis or alteration in red cell membrane permeability (5). Nakao et al (26) and Haradin et al (27) have demonstrated a progressive alteration in the shape of the stored red cells from a biconcave disc when fresh to a microspherocyte after 8 weeks. The progressive loss in total erythrocyte lipids accompanying in vitro storage in ACD which approximates 20 to 30% as 6 weeks (27) had been shown to be associated with losses of portions of the red cell membrane varying in size from macromolecular aggregates to microscopically visible buds (28, 29). Recent evidences indicate that the plasma  $K^+$

concentration increases in direct proportion to the storage time (25) and independent to small degree of hemolysis that occur during storage (7). At  $4^\circ C$  the rate of consumption of dextrose is approximately 0.05 mMoles/L. of red cells/hours (30, 31) which is at least 30 times less than at  $37^\circ C$  and the active transport of  $Na^+$  and  $K^+$  is almost halted. The increment plasma  $K^+$  concentration represents  $K^+$  that leaked from intact red cells, and the rate of efflux is inversely related to the ATP content of the stored red cells (32).

It is now realized that the deterioration of red cells during storage is a different process from aging in vivo. The ATP content of stored red

**Table 1.** Sequential Changes of Cellular Components of ACD Blood

Storage Time (days)	TOTAL WBC	WBC VIABILITY %	PMNs %	PLATELETS $\times 10^3 / \text{cu. mm.}$
0	11,925 $\pm$ 2,266	100	57.0 $\pm$ 7.17	309.0 $\pm$ 60.78
1	11,795 $\pm$ 3,265	99.7 $\pm$ 0.48	70.0 $\pm$ 7.12	289.8 $\pm$ 68.32
2	11,215 $\pm$ 2,412	96.9 $\pm$ 1.20	58.4 $\pm$ 9.96	272.2 $\pm$ 68.02
3	10,760 $\pm$ 2,496	91.8 $\pm$ 1.93	59.1 $\pm$ 13.67	252.1 $\pm$ 52.96
4	9,650 $\pm$ 2,158	88.6 $\pm$ 2.80	55.7 $\pm$ 10.27	243.8 $\pm$ 44.96
5	10,894 $\pm$ 2,282	84.5 $\pm$ 2.37	53.3 $\pm$ 13.90	224.6 $\pm$ 33.18
6	10,190 $\pm$ 2,183	81.0 $\pm$ 5.14	50.2 $\pm$ 18.15	226.5 $\pm$ 50.25
9	11,225 $\pm$ 3,464	72.1 $\pm$ 2.56	47.7 $\pm$ 17.67	205.6 $\pm$ 48.51
13	7,915 $\pm$ 1,720	66.0 $\pm$ 3.13	37.6 $\pm$ 14.66	143.1 $\pm$ 50.86

cells diminishes with time (25) but it undergoes little change while red cells age in vivo (33). Red cells lose their ability to survive in vivo before they hemolyzed in vitro. The damages in vitro which lead to loss of viability of red cells are not well understood. Factors which affect the osmotic fragility of stored red cells do not have any influence on their viability and substances which retard the rate

of hemolysis of stored blood do not necessarily prolong the viability of red cells (34). CPD solution is superior to ACD solution because it is more isotonic, better conservation of 2, 3, DPG and pCO<sub>2</sub>, maintains higher pH throughout storage with significantly lower plasma K<sup>+</sup> and free hemoglobin levels (4, 5) which should prolong the usable span of stored red cells. But there was little






FIGURE 1: The sequential changes of leukocytes, platelets, plasma potassium level and coagulation factors of ACD blood during storage.

difference in red cell survival when blood was stored in these two solutions up to 21 days (5). When blood was preserved for 2 to 3 weeks approximately 15 to 25% of transfused red cells were destroyed in vivo in the first 24 hours after transfusion (7). Thus the total  $K^+$  load from a transfusion can not be evaluated from plasma values alone. The red cell lysis in vivo must also be considered since its contribution may be nearly one-third of the total  $K^+$  load (7). Prolongation of the storage time by CPD results in high plasma levels of  $K^+$  ammonia and free hemoglobin (in addition to high  $Na^+$  concentration) (5). These substances could result in potentially deleterious effects in patients with hepatic and renal diseases.

The fall of total leukocyte count in our ACD blood to approximately 80% of the original value at 10 days of storage was comparable to that of the CPD blood (5, 6). The higher recovery of leukocytes in ACD and CPD blood than those observed in heparinized blood was probably related to the inhibition of surface adhesion of granulocytes in solution of low pH (35, 36) and decreased ionized calcium (37). The number of platelets in our ACD blood decreased to 61% by day 10 and to 46% at 2 weeks was slightly higher than those

observed in CPD blood (5). The decreased number of platelets and leukocytes observed is explained by the fact that they form clumps during storage (38-41). Light and electron microscopic examinations revealed that the microaggregates which developed in stored blood consist of degenerated platelets and leukocytes (40-42). Coincident with this there was a drop in the platelet count during the first week of storage and a progressive reduction in the absolute granulocyte count. The finding of higher recovery of platelets in our ACD blood is explained by the observations that platelets spontaneously clump more rapidly in CPD blood (8,9). This may partly be due to the effect of higher pH of CPD blood on the velocity of platelet aggregation (43). Lymphocytes are relatively nonadhesive (35) and maintain their functional and structural integrity during most of 3 weeks storage in ACD (44, 45). In contrast, platelets and granulocytes became adhesive and rapidly lose their viability during storage (35, 45, 46). Even though 66% of granulocytes were viable at 14 days of storage as observed by us it was demonstrated that irreversible loss of function of granulocytes occurred after a few hours and granulocyte mobility, phagocytosis and  $O_2$  consumption were markedly diminished.

after one week of storage in ACD plasma (18, 47). More recent observations indicated that granulocytes in whole blood stored for 24 hours retain their phagocytic and bactericidal activities (18, 45, 48). The decline in function after 24 hours of storage further indicated that the granulocytes would have only limited value for transfusion therapy thereafter. Platelets lose their ability to survive in vivo very rapidly and whole blood stored in ACD even 24 hours is a poor source of viable platelets (49, 51). It takes only 24 hours storage for the percentage of viable platelets to fall to 5% level (22).

Factor VIII activity dropped rapidly during storage at 4°C (52-54). Our baseline factor VIII activity was very low due to the dilution effect of large amount of ACD solution and delayed factor VIII determination. At 24 hours of collection the factor VIII activity was approximately 32% which is comparable to those observed by others (54, 55). Our results confirmed the instability of factor VIII and supported the recommendation that cryoprecipitate preparation should be processed within 4 hours of collection. The other coagulation factors known to deteriorate appreciably on storage is factor V. Factor V activity decreased rapidly to 40% of the initial concentration at day 7 comparable to

those observed by others (5, 55-57). Factor VII activity increased during storage similar to that observed earlier (55). This phenomenon is explained on the fact that cold-promoted enhancement of factor VII activity occurred by surface contact via factor XII and plasma kallikrein (58-62).

### บทคัดย่อ

คณะผู้รายงานได้ศึกษาการเปลี่ยนแปลงของส่วนประกอบต่างๆของเลือดจากคนธรรมดาจำนวน 20 ราย ที่เก็บไว้ใน ACD solution ที่ 4°C ในธนาคารเลือด ผลการศึกษาแสดงว่าเลือดที่เก็บไว้ใน ACD ในระยะสั้นคือประมาณ 2 สัปดาห์ มีคุณภาพเท่าเทียมกับที่เก็บไว้ใน CPD solution หลัง 24 ชั่วโมงแรก PMNs จะแตกทำลายไป ทำให้จำนวนของเม็ดเลือดขาวลดลงตามลำดับ เหลือ  $85.45 \pm 18.30\%$  ในสัปดาห์แรก และเหลือ 80% ในวันที่ 10 จำนวนของเกล็ดเลือดลดลงตามลำดับ เช่นเดียวกัน จนเหลือเพียง 46% ของค่าเดิมใน 2 สัปดาห์ ค่าที่ได้นั้นสูงกว่าเลือดที่เก็บไว้ใน CPD เนื่องจากเกล็ดเลือดเกาะกลุ่มกันน้อยกว่าที่เกิดขึ้นใน CPD ระดับของ factor VIII ลดลงอย่างรวดเร็วภายในเวลาไม่กี่ชั่วโมง ทำให้มีระดับเพียง 40% ภายในวันแรก แต่วันต่อไปก็มีระดับค่อนข้างคงที่ ระดับ factor V ลดลงช้าๆจนเหลือเพียง 40% ของค่าเดิมในวันที่ 7 ระดับ coagulation factor อื่นๆค่อนข้างคงที่ แต่ค่าของ factor VII กลับเพิ่มสูงขึ้นหลายเท่าตัว ภายในสัปดาห์แรก ทั้งนี้เป็นเพราะในขณะที่เก็บไว้ในความเย็นฤทธิ์ของ factor VII จะถูกกระตุ้นด้วย surface contact ผ่านทางระบบ Kallikrein.

## REFERENCES.

1. Lewisohn, R. : Blood transfusion by the citrate method. *Surg. Gynecol. Obstet.* 21:37, 1915.
2. Robertson O.H. : Transfusion with preserved red blood cells. *Brit. Med. J.* 1:691, 1918.
3. Loutit, J.F., Mollison, P.L., and Young, F.M. : Citrate acid-sodium citrateglucose mixtures for blood storage. *Q.J. Exp. Physiol.* 32:183 1943.
4. Gibson, G.J., II., Rees, S.B., McManus, T.J., and Scheitlin, W.A. : A citratephosphate-dextrose solution for the preservation of human blood. *Amer. J. Clin. Phat.* 28:569, 1957.
5. Limbird, T.J., and Silver, D. : Sequential changes in blood preserved with citrate-phosphate-dextrose. *Surg. Gynec. Obstet.* 138:401, 1974.
6. Bailey, D.N., and Bove, J.R. : Chemical and hematological changes in stored blood. *Transfusion* 15:244, 1975.
7. Simon, G.E., and Bove, J.R. : The potassium load from blood transfusion. *Postgrad. Med.* 49:61 1971.
8. Solis, R.T., Goldfinger, D., Gibbs, M.B., and Zeller, J.A. : Physical characteristics of microaggregates in stored blood. *Transfusion* 14: 538: 1974.
9. Becker, G.A., Chalos, M.K., Tuccelli, M., and Aster, R.H. : Prostaglandin F<sub>1</sub> in preparation and storage of platelet concentrates. *Science* 175:538, 1972.
10. Standard for Blood Banks and Transfusion Services. Fifth edition. Prepared by the Committee on Standards of the American Association of Blood Banks. Chicago. Twentieth Century Press, Inc., 1970.
11. Willis, J.B. : The determination of metals in blood serum by atomic absorption spectroscopy III, Sodium and potassium. *Spectrochim Acta* 16:551, 1960.
12. Lewis, M.L., and Ware, A.G. : A one-stage method for the determination of Accelerator Globulin. *Proc. Soc. Exp. Biol. Med.* 84:640, 1953.
13. Adamis, D., Sise, H.S., and Kimball, D.M. : The proconvertin test: a simplified method and its application to the study of anticoagulant processes. *J. Lab. Clin. Med.* 47:320, 1956.
14. Langdell, R.D., Wagner, R.H., and Brinkhous, K.M. : Effect of antihemophilic factor on one-stage clotting tests. *J. Lab. Clin. Med.* 41:637, 1953.
15. Sirridge, M.S. : Laboratory Evaluation of Hemostasis. Lea and Febiger. Phila., 1967.

16. Bachmann, F., Duckert, F., and Koller, F.: The Stuart-Prower factor assay and its clinical significance. *Thromb. Diath. Haemorrh.* 2:24, 1958.
17. Ratnoff, O.D., and Davie, E.W.: The purification of activated Hageman factor (activated factor XII). *Biochem.* 1:967, 1962.
18. Crowley, J.P., and Valeri, C.R.: Recovery and function of granulocytes stored in plasma at 4C for one week. *Transfusion* 14:574, 1974.
19. Michael, J.M., Doerner, I., Bruns, D., Ladenson, J.H., and Sherman, L.A.: Potassium load in CPD-preserved whole blood and two types of packed red blood cells. *Transfusion* 15:144, 1975.
20. Gibson, J.G., II, Gregory, C.B., and Button, L.N.: Citrate-phosphate-dextrose solution for preservation of human blood: a further report. *Transfusion* 1:280, 1961.
21. McCullough, J., and Weiblen, B.J.: Citrate phosphate dextrose (CPD) anticoagulant in blood transfusion. *Minn. Med.* 56:980, 1973.
22. Mollison, P.L.: *Blood Transfusion in Clinical Medicine.* F.A. Davis Co., Phila., 1967.
23. Orlina, A.R., and Josephson, A.M.: Comparative viability of blood stored in ACD and CPD. *Transfusion* 9:62, 1969.
24. Schechter, D.C., and Swan, H.: Biochemical alterations of preserved blood. *Arch. Surg.* 84:269, 1962.
25. Rapaport, S.: Dimensional, osmotic, and chemical changes of erythrocytes in stored blood. I. Blood preserved in sodium citrate, neutral and acid citrateglucose (ACD) mixtures. *J. Clin. Invest.* 26:591, 1947.
26. Nakao, M.T., Nakao, T., Tatibana, M., Yoshikawa, H., and Abe, T.: Effect of inosine and adenine on adenosine triphosphate regeneration and shape transformation in long-stored erythrocytes. *Biochem. Biophys. Acta* 32:564, 1959.
27. Haradin, A.R., Weed, R.I., and Reed, C.F.: Changes in physical properties of stored erythrocytes. Relationship to survival in vivo. *Transfusion* 9:229, 1969.
28. Reed, C.F., Swisher, S.N., Marinetti, G.V., and Eden, E.G.: Studies on the lipids of the erythrocytes. I. Quantitative analysis of the lipids of normal human red blood cells. *J. Lab. Clin. Med.* 56:281, 1960.
29. Weed, R.I., and Bowdler, A.J.: Metabolic dependence of the critical hemolytic volume of human erythrocytes: relationship to osmotic fragility and autohemolysis in hereditary spherocytosis and normal red blood cells. *J. Clin. Invest.* 45:1137, 1966.

30. Strumia, M.M.: Methods of blood preservation in general and preparation and use of red cell suspensions. *Amer. J. Clin. Path.* 24:260, 1954.
31. Chaplin, H. Jr., Crawford H., Cutbush, M., and Mollison, P.L.: Posttransfusion survival of red cells stored at -20°C. *Lancet* 1:852, 1954.
32. Dern, R.J., Brewer, G.J., and Wiorowski, J.J.: Studies on the preservation of human blood. II. The relationship of erythrocyte adenosine triphosphate levels and other in vitro measures to red cell storage-ability. *J. Lab. Clin. Med.* 69:968, 1967.
33. Gabrio, B.W. and Finch, C.A.: Erythrocyte preservation. I. The relation of the storage lesion to in vivo erythrocyte senescence. *J. Clin. Invest.* 33:242, 1954.
34. Chaplin, H., Jr., Crawford, H., Cutbush, M., and Mollison, P.L.: The effects of a phenothiazine derivative (RP 3300) on red cell preservation. *J. Clin. Path.* 5:91, 1952.
35. Garvin, J.E.: Factors affecting the adhesiveness of human leukocytes and platelets in vitro. *J. Exp. Med.* 114:51, 1961.
36. Lycette, R.M.: Influence of ACD and CPD anticoagulants on white cell aggregation. *Transfusion* 13: 435, 1973.
37. Moore, E.W.: Ionized calcium in normal serum, ultrafiltrates and whole blood determined by ion-exchange electrodes. *J. Clin. Invest.* 49:318, 1970.
38. Tullis, J.L.: Preservation of leukocytes. *Blood* 8:563, 1953.
39. Kattlove, H.E., and Alexander, B.: The effect of cold on platelets. I. Cold-induced platelet aggregation. *Blood*. 33:39, 1971.
40. Swank, R.L.: Alterations of blood on storage: Measurement of adhesiveness of "aging" platelets and leukocytes and their removal by filtration. *N. Eng. J. Med.* 265:728, 1961.
41. Kartasheosky, N.G. and Rumyantsev, V.V.: Microclots of stabilized blood. *Probl. Gematol. Pareliv. Krovi.* 13:6, 1968.
42. Zucker, M.B., and Borelli, J.: Viscous metamorphosis produced by chilling and by clotting. Failure to find specific defect of viscous metamorphosis in PTA syndrome. *Thromb. Diath. Haemorrh.* 4:424, 1960.
43. Skoza, L., Zucker, M.B., Jerushalmy Z., and Grant, R.: Kinetic studies of platelet aggregation induced by adenosine diphosphate and its inhibition by chelating agents, guanidino compounds and adenosine. *Thromb. Diath. Haemorrh.* 18:713, 1967.

44. Petrakis, N.L., and Politis, G.: Prolonged survival of viable mitotically competent mononuclear leukocytes in stored whole blood. *N. Eng. J. Med.* 267:286, 1962.
45. McCullough, J., Benson, S.J., Yunis, E.J., and Quie, P.G.: Effect of blood-bank storage on leukocyte function. *Lancet* 2:1333, 1969.
46. Jenevein, E.P., and Weiss, D.L.: Platelet microemboli associated with massive blood transfusion. *Amer. J. Pathol.* 45:313, 1964.
47. Strumia, M.M.: Preservation of leukocytes. In the "Preservation of the Formed Elements and of the Proteins of the Blood. American National Red Cross. 1949.
48. McCullough, J., Carter, S.J., and Quie, P.G.: Effects of anticoagulants and storage on granulocyte function in bank blood. *Blood.* 43:207, 1974.
49. Baldini, M., Costea, N., and Dameshek, W.: The viability of stored human platelets. *Blood.* 16:1669, 1960.
50. Morrison, F.S., and Baldini, M.: The favorable effect of ACD on the viability of fresh and stored human platelets. *Vox. Sang.* 12 : 90, 1967.
51. Levin, R.H., Freireich, E.J., and Chappell, W.: Effect of storage up to 48 hours on response to transfusion of platelet rich plasma. *Transfusion* 4:251, 1964.
52. Penick, G.D., and Brinkhous K.M.: Relative stability of plasma antihemophilic factor (AHF) under different conditions of storage. *Amer. J. Med. Sci.* 232 : 434, 1956.
53. Bowie, E.J., Thompson, J.H., and Owens, C.A.: The stability of antihemophilic globulin and labile factor in human blood. *Mayo Clinic Proc.* 39:144, 1964.
54. Schlichter, S.J., Counts, R.B., Henderson, R., and Harker, L.A.: Preparation of cryoprecipitated factor VIII concentrates. *Transfusion* 16:616, 1976.
55. Stefanini, M., and Dameshek W.: *The Hemorrhagic Disorders.* Second Edition. Grune and Stratton, New York, 1962.
56. Goldstein, R., Bunker J.P., and McGovern, J.J.: The effect of storage of whole blood and anticoagulants upon certain coagulation factors. *Ann. N.Y. Acad. Sci.* 115:422, 1964.
57. Mooreside, D.E., Graybeal, F.Q., Jr., and Langedell, R.D.: Effects of adenine on clotting factors in fresh blood, stored blood, and stored fresh frozen plasma. *Transfusion* 9:191, 1969.
58. Rapaport, S.I., Aas, K., and Owren, P.A.: The effect of glass upon the activity of the various plasma clotting factors. *J. Clin. Invest.* 34:9, 1955.

59. Altman, R., and Hemker, H.C.: Contact activation of the extrinsic blood clotting system. Thromb. Diath. Haemorrh. 18:525, 1967.
60. Gjonnaess, H.: Cold - promoted activation of factor VII. III. Relation to the kallikrein system. Thromb. Diath. Haemorrh. 28: 182, 1972.
61. Gjonnaess, H.: Cold - promoted activation of factor VII. IV. Relation to the coagulation system. Thromb. Diath. Haemorrh. 28: 194, 1972.
62. Saito, H., and Ratnoff, O.D.: Alteration of factor VII activity by activated Fletcher factor (a plasma kallikrein): a potential link between the intrinsic and extrinsic blood-clotting systems. J. Lab. Clin. Med. 85:405, 1975.

## ห้างฯ เชียงใหม่วินเนอร์

๙๕/๓ ถนนห้วยแก้ว เชียงใหม่

จำหน่าย เครื่องมือเครื่องใช้ในการแพทย์  
เครื่องมือวิทยาศาสตร์ เครื่องแก้ว สารเคมี ไซริงค์  
และเข็มฉีดยา น้ำยาตรวจวิเคราะห์โรค อาหาร  
เลี้ยงเชื้อ ฯลฯ

บริการรวดเร็ว และราคาเบา