

Study of Stress Experienced by Parents about Their Children and Adolescents during COVID-19 Pandemic. Fayoum Governorate. Egypt

Remon M. Yousef¹, Shaimaa S. Soliman², Mariam E. Dawoud³

¹Pediatrics Department, Faculty of Medicine, Fayoum University, Egypt.

²Public health and community medicine Department, Faculty of Medicine, Menoufia University, Egypt.

³Psychiatry Department, Faculty of Medicine, Fayoum University, Egypt.

How to Cite: Remon M. Y., Shaimaa S. S., Mariam E. D. (2022). Study of Stress Experienced by Parents about Their Children and Adolescents during COVID-19 Pandemic. Fayoum Governorate. Egypt. *International Journal of Child Development and Mental Health*, 10(2), 11-18.

***Corresponding author:**

Email: med11@fayoum.edu.eg
Tel. 002-01200611326

Article Info:

Received: 5 March 2022

1st Revision: 14 September 2022

Accepted: 5 October 2022

Keywords:

Coping, COVID, Parents, Stress

Abstract

a major external stressor, COVID-19 pandemic with all its negative consequences as the lockdown, social obligations, changes in routine and online schooling, health concerns, rapidly increasing affected cases and deaths constitutes a major health concern which heavily affected the psychological wellbeing. Parents are highly vulnerable group and showed high levels of perceived stress during this critical period. This study aimed to assess the stress experienced by parents during the COVID period. Online Google forms and hard copies contained demographic data, perceived stress scale and risk factors were filled by parents (with age ranges from 20-60) of offspring aged till 18 years old. 1265 valid questionnaires with vast majority (93.9%) were mothers. Moderate degree of stress was found in 70.3%. Multivariate regression model for predicting PSS was significantly related to the age category, parenthood and history of psychic trauma. Majority of parents experienced moderate to high grades of stress. The perceived stress was more evident in mothers, also affected by other factors which affect resilience like age, history of previous psychic trauma , and history of COVID infection.

Introduction

The COVID-19 pandemic is increasing rapidly; world health organization reported 5,819,962 confirmed cases worldwide in May, 2020, while in Egypt 22,082 confirmed cases was reported and 879 deaths have been in May 2020. On 24th March 2021 there have been 124,535,520 confirmed cases of COVID-19, including 2,738,876 deaths, and in Egypt 197,350 confirmed cases of COVID-19 with 11,720 deaths (WHO, 2020) Egypt started restrictive measures and quarantine, with restriction regarding school attendance to avoid the risk of spread of infection since mid of March 2020.

Epidemiological data suggest that children have the same possibility to catch COVID as adults yet, milder (Qiu et al., 2020). Data available at the time of the publication of Qiu et al. 2020 suggested that children account for between one and 5% of COVID-19 cases. Those aged less than 1 year appear most likely to suffer critical infection (Covid et al., 2020; Centers for Disease Control and Prevention, 2020; Dong et al., 2020). Deaths in children are extremely rare. However, there are increasing reports of complications among children and adolescents (Castagnoli et al., 2020) at this points not only children are stressed but also their parents experience

high level of stress. Also, COVID 19 outbreak and pandemic has great challenges with disruption of daily life routine, financial losses, restrictions about isolation at home, sanitation and obligation about social distancing and infection control, confusion about school opening or not and uncertainty about actual management protocol or effective vaccine which may constitute potential for increased family stress. Moreover, the lockdown in general meant that children were unable to access leisure activities, clubs...etc. all of which affected stress perception among children and their parents. (Leigh-Hunt et al., 2017)

Objectives

To study the stress among parents during COVID-19 pandemic and worry experienced regardless the children' age or gender and find the relation between stress and other risk factors.

Study design

A cross-sectional survey of stress during Covid-19 pandemic in Egypt that target on parents' response.

Methods

The author designed Online Google forms and hard copies which contain demographic data, perceived stress scale, risk factors which may increase the perceived stress during COVID 19 outbreak. The online forms were launched through social media also, hard copies were designed to be filled by parents who are attendants to pediatric clinic and psychiatry clinic, Fayoum University Hospitals with a total of 1265 valid forms collected. All parents who have children (0-18years) were eligible to fill the form. Both forms were available to be filled by the participants through the period from first of October 2020 till first of November 2020. The designed period was associated with the decision of the Egyptian ministry of Education about return to nurseries, kindergarten, schools, and universities. Informed consent was included at the beginning of the form and approval is mandatory to fill the form. The questionnaires were anonymous to ensure the confidentiality and reliability of the data.

The study and the consent were approved by the ethical committee, Faculty of Medicine Fayoum University, (no. 126). Forms contained demographic data (Parents' age, whether father or mother, and their jobs), past history of experiencing life trauma (loss, accidents or abuse), history of COVID 19 infection, multiple choice question about the most suggested causes of stress during this period, items of the perceived stress scale and open-ended question inquiring the parents' point of view about how to accept and overcome their perceived stress.

Psychometric tool

The Perceived Stress Scale (PSS) (Cohen, Kamarck, & Mermelstein, 1983) is a tool for measuring psychological stress during the previous month. It is a 10-item scale in which each item has graded from 0 to 4, 0 means never experienced while 4 means very often experienced. We used the scale in Arabic version; this was validated in Arabic among pregnant and postpartum women (Chaaya, Osman, Naassan, & Mahfoud, 2010) Scores ranging from 0-13 would be considered low stress, scores ranging from 14-26 would be considered moderate stress, and scores ranging from 27-40 would be considered high perceived stress.

Statistical analysis of the collected data

An IBM compatible personal computer with SPSS statistical package version 23 was used to analyze Data (SPSS Inc., 2015. International Business Machines Corporation (IBM) SPSS statistics for windows, Armnok, NY, version 23.0). The variables were expressed in: Number (No), percentage (%) mean (\bar{x}) and standard deviation (SD). Association between qualitative variables was assessed using Chi-square test (χ^2). Fischer's Exact test was used in the case that any of the expected cells were less than five. A logistic regression was performed to ascertain the effects of possible risk factors on depression, anxiety and other outcomes. Two-sided P value <0.05 was considered statistically significant.

Results

The study included 1265 valid questionnaires. About two thirds (70.6%) of the participants were in the age

group of 31-40 years and their vast majority (93.9%) were mothers. More than half of them (53.1%) did not

have any history of psychic trauma and only 5.1% had history of COVID-19 infection (Table 1).

Table 1: participants' characteristics: (Arrange a new table as a standard format)

| Character | No. (%) |
|----------------------------------|-------------|
| Age category | |
| 20-30 | 255 (20.2) |
| 31-40 | 893 (70.6) |
| 41-50 | 110 (8.7) |
| 51-60 | 7 (0.6) |
| Parenthood | |
| Mother | 1188 (93.9) |
| Father | 77 (6.1) |
| History of psychic trauma | |
| Absent | 672 (53.1) |
| Present | 593 (46.9) |
| COVID history | |
| Absent | 1191 (94.2) |
| Present | 65 (5.1) |

The mean score of the PSS was 21.79 (ranging from 4.0 to 36.0). Most of the participants (70.3%) had a

moderate grade of stress, while 21.1% had the high grade and only 8.6% had low grade of stress (Table 2)

Table 2: PSS mean score and grades.

| Variable | Mean \pm SD, range |
|------------|----------------------------|
| PSS | 21.79 \pm 5.67, 4.0-36.0 |
| PSS grades | No. (%) |
| Low | 109 (8.6) |
| Moderate | 889 (70.3) |
| High grade | 267 (21.1) |

The mean PSS varied significantly among different age groups ($P < 0.001$), as it was significantly lower in the age group of 41-50 years than the mean PSS in both age groups of 20-30 y. and 31-40 years ($p < 0.001$

and 0.002 respectively) The mean PSS was significantly higher among mothers than fathers ($p < 0.001$), among participants with history of psychic trauma ($p < 0.001$) or history of COVID -19 infection (0.041) (Table 3).

Table 3: mean PSS scores and possible risk factors:

| Risk factors | PSS (mean) | PSS (SD) | P value |
|---------------------------|------------|----------|---------|
| Age category | | | |
| 20-30 | 22.53 | 5.45 | 0.001 |
| 31-40 | 21.81 | 5.63 | 0.002 |
| 41-50 | 20.00 | 6.12* | |
| 51-60 | 19.71 | 7.01 | |
| Parenthood | | | |
| Mother | 21.94 | 5.62 | <0.001 |
| Father | 19.46 | 6.01 | |
| History of psychic trauma | | | |
| Absent | 21.15 | 5.80 | <0.001 |
| Present | 22.51 | 5.43 | |
| COVID-19 history | | | |
| Absent | 21.74 | 5.68 | 0.041 |
| Present | 23.21 | 5.32 | |

* The PSS in the age group of 41-50 years was significantly lower than the PSS in the age group 20-30 years ($P < 0.001$) and the age group 31-40 years ($p=0.002$)

Among all age categories, moderate grades of PSS was most prevalent, this was followed by the high grades except for the age category of 41-50 years ($p = 0.007$). The low grade of PSS was lowest in prevalence. This difference in prevalence was significant among age category of 20-30 and 41-50 years (Table 4) In any of the parents, low grades of PSS

were significantly lower than moderate or high grades ($P = 0.005$). Among participants with and without history of psychic trauma, moderate grades of PSS were significantly higher and low grades of PSS were significantly lower than other grades ($p = 0.001$). History of COVID-19 infection was not significantly related to the grade of PSS ($P = 0.179$) (Table 4).

Table 2: ADHD Score of SNAP-iv scale by parents and teachers.

| Risk factors | PSS grade | | | P value |
|---------------------------|------------------------|-----------------------------|-------------------------|---------|
| | Low (n=109) No. (%) | Moderate (n=889) No. (%) | High (n=267) No. (%) | |
| Age category | | | | |
| 20-30 | 17 (6.7) | 171 (67.1) | 67 (26.3)* | 0.007 |
| 31-40 | 73 (8.2) | 636 (71.2) | 184 (20.6) | |
| 41-50 | 18 (16.4) † | 78 (70.9) | 14 (12.7) | |
| 51-60 | 1 (14.3) | 4 (57.1) | 2 (28.6) | |
| Parenthood | | | | |
| Mother | 95 (8.0) † | 837 (70.5) | 256 (21.5) | 0.005 |
| Father | 14 (18.2) † | 52 (67.5) | 11 (14.3) | |
| History of psychic trauma | | | | |
| Absent | 74 (11.0) † | 475 (70.7)* | 123 (18.3) | 0.001 |
| Present | 35 (5.9) † | 414 (69.8)* | 144 (24.3) | |
| COVID-19 history | | | | |
| Absent | 104 (8.7) | 839 (70.4) | 248 (20.8) | 0.179 |
| Present | 3 (4.6) | 43 (66.2) | 19 (29.2) | |

† Significantly lower, * significantly higher than their corresponding in the other categories

Linear regression model for predicting PSS was significantly related to the age category, parenthood, history of psychic trauma and history of COVID-19 infection ($p < 0.001$, <0.001 , <0.001 and 0.041

respectively). By multivariate regression, all of these factors remained significant except COVID-19 infection history ($p < 0.001$, 0.001 , <0.001 and 0.102 respectively) (Table 5)

Table 5: Univariate and multivariate regression analysis of possible risk factors.

| Risk factors | Univariate | | | Multivariate | | |
|---------------------------|------------------|----------------|---------|------------------|----------------|---------|
| | Unstandardized B | Standardized B | P value | Unstandardized B | Standardized B | P value |
| Age category | -1.095 | -0.106 | <0.001 | -1.029 | -0.099 | <0.001 |
| Parenthood | -2.474 | -0.104 | <0.001 | -2.243 | -0.095 | 0.001 |
| History of psychic trauma | 1.366 | 0.120 | <0.001 | 1.362 | 0.120 | <0.001 |
| COVID history | 1.474 | 0.058 | 0.041 | 1.169 | 0.046 | 0.102 |

Discussion

This study reflected the increased degree of distress among parents during the period of COVID pandemic. The majority of parents experienced

moderate degrees of stress, which increased more among those with past history of psychic trauma and history of COVID infection. A review of literature revealed that high stress in general is a common reaction to pandemics also; stress is common among

parents than non-parents (American Psychological Association, 2020), though the nature of the current pandemic is totally different with the great restrictive measures, social distancing, wearing masks, school closure, rapid deaths, rapid mutation, vague nature.. etc. Our work revealed that mothers were significantly higher in perceiving stress as previously reported by Bjørknes et al. (2022) in one year follow up study; sensitive and caring nature of mothers may support this result. Contrary to authors' expectations, the age group (41-50 years) was the highest in stress. The point is this age group may have more coping strategies, more financial stability. However, participants of this age in Egypt may have grandchildren not only children which may raise fear about the pandemic.

However multiple factors may contribute to increased level of stress among parents during COVID stress, a study of Norwegian parents found that mothers with more than one child and parents with a psychiatric diagnosis reported greater levels of parental stress, more burnout, and more anger towards their children, as well as less social support (Johnson, Skjerdingstad, Ebrahimi, Hoffart, & Johnson, 2022). In our study, Linear regression model for predicting PSS was significantly related to history of psychic trauma and history of COVID-19 infection and in accordance with this Marzilli et al. (2021) found that Parents' past traumas significantly predicted peritraumatic distress due to COVID-19 and children's psychological difficulties. Research by Demir et al. (2020) has shown that having experienced previous trauma may raise the possibility to the adverse effects of subsequent traumatic experiences when the prior traumatic exposure sensitizes the autonomic nervous system with subsequent adverse mental and physical effects. History of COVID infection may booster the adverse mental effect due to the acknowledgment of being in danger which may constitutes constitute a traumatic experience itself. However, survivors of this infection at the time of pandemic were expected to gain more resilience. Cusinato et al. (2020) showed that certain variables regarding socio-demography affected parental stress, changes in working conditions, younger children, and parental psychological, physical, or genetic problems appeared

to be linked to lower levels of psychological well-being in parents during the COVID-19 lockdown. Family structure, the number, and the presence of children with psychological, physical, or genetic diseases are all risk factors for high parental stress. As a response to a multiple choice question concerning causes of stress which may be experienced by parents in the period of the pandemic, the most suggested causes were worry about infection of their children during the school attendance, uncertainty about the available protocol of management of COVID, the uncertainty about the efficacy of the available vaccines, and the affection of their offspring with psychiatric disorders as a result of the stress during the period of COVID pandemic. While (Fiese et al., 2002) reported a change in children's daily structure and routines as the most stressful factors for parents during the pandemics as it makes children feel more insecure with more problematic behavior. Other reports found that online learning and deprivation from friends were important stressors for children and their parents during the pandemic (Covid et al., 2020; Centers for Disease Control and Prevention, 2020). Hearing and reading about the severity and contagiousness of COVID were found to be another important stressor which may increase the negative psychological impact among parents (Park et al., 2020; Adams, Smith, Caccavale, & Bean, 2021).

In this study parents reported that praying with their children, infection control measures, and staying at home were the most used coping strategy among them in a trial to reduce their stress and anxiety about the pandemic. Achterberg, Dobbelaar, Boer, & Crone (2021) studied the longitudinal effect of COVID related stress on parents and children wellbeing and found that online schooling, working remotely or being unable to work at all, financial and health concerns for the family were risk factors related to increased depression, anxiety, hostility and interpersonal sensitivity among both parents and children. Also, the previously mentioned factors affected the parents and children bonding with increased risk of internalizing and externalizing disorders among children. The aforementioned study found that the poor family wellbeing prior COVID stress and poor coping (parental over reactivity)

during COVID period were related to poor coping. High levels of positive parental coping (positive reappraisal) and reflection could have positive consequences by increasing family bonding and improve parent-child relations Achterberg et al. (2021). In accordance with our results Brown, Doom, Lechuga-Peña, Watamura, & Koppels. (2020) studied stress and parenting during COVID pandemic among 183 participants who were parents of children under 18 years old, in which mothers were 89.6 % and fathers were 10.4% and found the mean and standard deviation of the perceived and stress scale was 19.14 (6.59), with range (0-37). The latter study found that most of parents experienced mood changes, anxiety, stress, and sleep problems and this was mostly because their children experienced changes in learning and health. Also, the author found the relation between the parental stress and increased child abuse Brown et al. (2020). Also, a Chinese study designed online questionnaire to measure students' parents stress during COVID outbreak with 1163 valid questionnaires collected, 80.2% were mothers and 19.8% were fathers (Zimmermann & Curtis, 2020). In the aforementioned study no gender difference was found in contrary to our results, this is may be due to low number of fathers participants in our representative sample. In our study stress level was significantly lower in parents aged 41-50 years old despite their offspring age or grade as our study didn't include the age of them while in a study by Wu et al. (2020), anxiety, depression and stress were significantly lower in parents of college students being more socially supportive to their parents.

Different studies addressed different factors related to stress perceived during COVID pandemic according to their studied variables and tested their correlations using regression analysis. In a Chinese study, which tested 1163 valid online questionnaire to study students' parents stress during COVID, in which regression analysis results showed that perceived stress, social support, marital satisfaction, family conflicts, child's learning stage as well as parents' history of mental illness had significant effects on parents' anxiety and depression (27) where social support and satisfying marriage were protective factors against stress. In a study on Italian sample

Giannotti, Mazzoni, Bentenuto, Venuti, & de Falco (2021) studied parental stress and externalizing behavior among children during the COVID lockdown to find that, in the final regression model, there was a significant effect of parent gender, child age, time dedicated by the parent doing activities with the child, and quality of co-parenting relationship. Also, Giannotti et al. (2021) found that distant learning and workload were significant predictors of parental stress in parents whom their children were involved in distant learning. Regarding risk predictors and protective factors of parental stress related to COVID among parents with children who have special conditions like CP, ADHD and other neurodevelopmental disabilities different studies were done. Montirossi et al. (2021) studied stress and resilience among 1,472 parents (83.1% mothers) of children and adolescents with neurodevelopmental disorders to find that stepwise regression analysis revealed that perception of self, planned future, family cohesion were protective factors buffering parental stress during COVID period.

Limitation of the study

The study wasn't limited to certain child age group also, the childrens' age wasn't included in the forms. Only one valid psychometric tool was used to measure the perceived stress which may not draw a well conclusive data. Response options for questions designed by the researchers may not have encompassed all possible stressors. Stress measured only once and wasn't reassessed again. Fathers as participants were much lower in number than mothers and this gender difference may constitute a confounding factor.

Conclusion

Majority of parents experienced moderate to high grades of stress. The perceived stress was more evident in mothers, also affected by other factors which affect resilience like age, past history of previous psychic trauma, history of COVID infection. This study revealed how far the mental health of parents could be affected hence lead to poor quality of life and poor parenting.

Recommendation

Evidence based resources for supporting parents are mandatory for providing them with information about culture-based stress coping skills. Already efforts were done from organization like center for disease control and prevention (CDC) and world health organization (WHO) to provide online coping with stress resources for parents (Parenting for Lifelong Health, 2020; World Health Organization. Looking after our mental health. 2020).

Declaration

Ethics approval and consent to participate:

This study was approved by the Research Ethics Committee, Faculty of Medicine, Fayom University. The number of approval is (r 126). Contents of consent were clarified and written at the start of the online questionnaire and approval was necessary to complete the form.

Consent for publication: Not applicable.

Availability of data and materials: Not applicable

Competing interests: "No competing interests were declared" in this section.

Funding: This study didn't receive any fund.

Authors' contributions

RY prepared the main idea, distributed the forms both online and in the clinics and contributed in writing the manuscript. MD prepared the Google form, analyzed and interpreted the patient data regarding the clinical data and psychometric tools and was a major contributor in writing the manuscript. SS was the major contributor in statistical analysis. All authors read and approved the final manuscript

Acknowledgments

The authors express their deep gratitude to all the participants.

Reference

Achterberg, M., Dobbelaar, S., Boer, O.D. & Crone, E.A. (2021). Perceived stress as mediator for longitudinal effects of the COVID-19 lockdown on wellbeing of parents and children. *Scientific reports*, 11(1), 1-14.

Adams, E. L., Smith, D., Caccavale, L. J., & Bean, M. K. (2021). Parents are stressed! Patterns of parent stress across COVID-19. *Frontiers in psychiatry*, 12, 626456.

American Psychological Association. (2020). *Stress in the time of COVID-19*. Volume one. Retrieved from: <https://www.apa.org/news/press/releases/stress/2020/report>.

Bjørknes, R., Skogen, J. C., Nærde, A., Sandal, G. M., Haug, E., Mæland, S., ... & Lehmann, S. (2022). Parental stress during the COVID-19 pandemic: A one-year follow-up. *Plos one*, 17(12), e0276190.

Brown, S. M., Doom, J. R., Lechuga-Peña, S., Watamura, S. E., & Koppels, T. (2020). Stress and parenting during the global COVID-19 pandemic. *Child abuse & neglect*, 110, 104699.

Castagnoli, R., Votto, M., Licari, A., Brambilla, I., Bruno, R., Perlini, S., ... & Marseglia, G. L. (2020). Severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infection in children and adolescents: a systematic review. *JAMA pediatrics*, 174(9), 882-889.

Covid, C. D. C., Team, R., COVID, C., Team, R., Bialek, S., Boundy, E., ... & Sauber-Schatz, E. (2020). Severe outcomes among patients with coronavirus disease 2019 (COVID-19)-United States, February 12-March 16, 2020. *Morbidity and mortality weekly report*, 69(12), 343-346.

Centers for Disease Control and Prevention. Coping with stress. *Coronavirus disease 2019 (COVID-19)*. National Center for Immunization and Respiratory Diseases, Division of Viral Diseases. (2020). Retrieved August 25, 2020, From <https://www.cdc.gov/coronavirus/2019-ncov/daily-life-coping/managing-stress-anxiety.html>.

Chaaya, M., Osman, H., Naassan, G., & Mahfoud, Z. (2010). Validation of the Arabic version of the Cohen Perceived Stress Scale (PSS-10) among pregnant and postpartum women. *BMC psychiatry*, 10(1), 1-7.

Cohen, S., Kamarck, T., & Mermelstein, R. (1983). A global measure of perceived stress. *Journal of health and social behavior*, 385-396.

Cusinato, M., Iannattone, S., Spoto, A., Poli, M., Moretti, C., Gatta, M. & Micsioscia, M. (2020). Stress, resilience, and well-being in Italian children and their parents during the COVID-19 pandemic. *International Journal of Environmental Research and Public Health*, 17(22), 8297.

Demir, Z., Böge, K., Fan, Y., Hartling, C., Harb, M. R., Hahn, E., Seybold, J., & Bajbouj, M. (2020). The role of emotion regulation as a mediator between early life stress and posttraumatic stress disorder, depression and

anxiety in Syrian refugees. *Translational Psychiatry*, 10(1), 1-10.

Dong, Y., Mo, X. I., Hu, Y., Qi, X., Jiang, F., Jiang, Z., & Tong, S. (2020). Epidemiological characteristics of 2143 pediatric patients with 2019 coronavirus disease in China. *Pediatrics*, 145(6), e20200702.

Fiese, B. H., Tomcho, T. J., Douglas, M., Josephs, K., Poltrack, S., & Baker, T. (2002). A review of 50 years of research on naturally occurring family routines and rituals: Cause for celebration?. *Journal of family psychology*, 16(4), 381.

Giannotti, M., Mazzoni, N., Bentenuto, A., Venuti, P., & de Falco, S. (2021). *Family adjustment to COVID-19 lockdown in Italy: Parental stress, coparenting, and child externalizing behavior*. Family Process.

Johnson, M. S., Skjerdingstad, N., Ebrahimi, O. V., Hoffart, A., & Johnson, S. U. (2022). Parenting in a Pandemic: Parental stress, anxiety and depression among parents during the government-initiated physical distancing measures following the first wave of COVID-19. *Stress and Health*, 38(4), 637-652.

Leigh-Hunt, N., Bagguley, D., Bash, K., Turner, V., Turnbull, S., Valtorta, N., & Caan, W. (2017). An overview of systematic reviews on the public health consequences of social isolation and loneliness. *Public health*, 152, 157-171.

MacKinnon, D. P., Fairchild, A. J., & Fritz, M. S. (2007). Mediation analysis. *Annual review of psychology*, 58, 593.

Marzilli, E., Cerniglia, L., Tambelli, R., Trombini, E., De Pascalis, L., Babore, A., Trumello, C., & Cimino, S. (2021). The COVID-19 Pandemic and Its Impact on Families' Mental Health: The Role Played by Parenting Stress, Parents' Past Trauma, and Resilience. *International journal of environmental research and public health*, 18(21), 11450.

Montiroso, R., Mascheroni, E., Guida, E., Piazza, C., Sali, M.E., Molteni, M. & Reni, G. (2021). Stress symptoms and resilience factors in children with neurodevelopmental disabilities and their parents during the COVID-19 pandemic. *Health Psychology*, 40 (7), 428.

Parenting for Lifelong Health. (2020). *COVID-19: 24/7 Parenting*. Retrieved August 25, 2020 from <https://www.covid19parenting.com>.

Park, C. L., Russell, B. S., Fendrich, M., Finkelstein-Fox, L., Hutchison, M., & Becker, J. (2020). Americans' COVID-19 stress, coping, and adherence to CDC guidelines. *Journal of general internal medicine*, 35(8), 2296-2303.

Qiu, H., Wu, J., Hong, L., Luo, Y., Song, Q., & Chen, D. (2020). Clinical and epidemiological features of 36 children with coronavirus disease 2019 (COVID-19) in Zhejiang, China: an observational cohort study. *The Lancet infectious diseases*, 20(6), 689-696.

Rajkumar, R. P. (2020). COVID-19 and mental health: A review of the existing literature. *Asian journal of psychiatry*, 52, 102066.

World Health Organization. (2020). *Coronavirus disease 2019 (COVID-19): situation report*, 73.

World Health Organization. *Looking after our mental health. #HealthyAtHome – Mental health*. Retrieved August 25, (2020). from <https://www.who.int/campaigns/connecting-the-world-to-combat-corona-virus/healthyathome/healthyathome---mental-health>.

Wu, M., Xu, W., Yao, Y., Zhang, L., Guo, L., Fan, J. & Chen, J. (2020). Mental health status of students' parents during COVID-19 pandemic and its influence factors. *General Psychiatry*, 33(4).

Zimmermann, P., & Curtis, N. (2020). Coronavirus infections in children including COVID-19: an overview of the epidemiology, clinical features, diagnosis, treatment and prevention options in children. *The Pediatric infectious disease journal*, 39(5), 355.