

# **Frontoethmoidal meningoencephalocele with multiple congenital craniofacial and skeletal malformations in a stillborn puppy**

**Promporn Raksaseri<sup>1</sup> Wuthichai Klomkleaw<sup>1\*</sup>**

## ***Abstract***

A one-and-a-half-year-old Shih-Tzu bitch went into natural labor after full term of her first pregnancy, giving birth to 4 puppies. One of them was a male which died at birth, 104 grams in weight and 12 centimeters in length. This puppy showed several facial abnormalities and skeletal deformities. It was preserved in 10% formalin for anatomical, histopathological, and radiographic studies. Gross anatomical study revealed congenital anomalies of the eyes including anophthalmia of the left eye and microphthalmia of the right eye. Malformations of the ears including anotia of the left ear and microtia of the right ear were observed. Evidence of cleft lips was also noted. There was a membrane-bound mass emerging from the center of the face and replacing the nose. Histopathological sections of the mass revealed evidence of nervous tissues composed of neurons and their supporting cells including microglia and astrocytes. Radiographs showed incomplete formation of frontal bones where the herniation of the mass occurred. These findings were consistent with a diagnosis of frontoethmoidal meningoencephalocele. Additionally, deformities of the 1<sup>st</sup> to the 3<sup>rd</sup> cervical vertebrae, the 3<sup>rd</sup> and lower thoracic vertebrae, and all lumbar vertebrae were found. Rib deformity was observed at both sides and rib fusion was presented on the right side. In conclusion, this study shows a rare case of a stillbirth puppy that possessed frontoethmoidal meningoencephalocele associated with multiple facial and skeletal abnormalities.

---

**Keywords:** frontoethmoidal encephalomeningocele, facial anomalies, skeletal abnormalities, inherited, dog

<sup>1</sup>Department of Anatomy, Faculty of Veterinary Science, Chulalongkorn University, Bangkok 10330, Thailand

\*Correspondence: Wuthichai.K@chula.ac.th

## Introduction

Congenital malformations or congenital anomalies are defect conditions at birth which can be found in many parts of the body. These conditions can be caused by toxic substances, drug toxicity, injury during pregnancy, teratogens, malnutrition of the bitch and genetic defects. In case of canine facial malformations, there are several congenital defects including cyclopion, cleft lip and cleft palate, and frontoethmoidal meningoencephalocele. One of the most common congenital defects of dog face is cleft lip, which might be associated with cleft palate resulting from malformation of the jaw and the face during embryogenesis.

Frontoethmoidal meningoencephalocele is a protrusion of intracranial tissues and meninges through the frontal bone (David, 1993; Suwanwela and Suwanwela, 1972). It can be congenital or acquired disorders caused by tumors, hydrocephalus, or other causes (Holmes et al., 2001). The brain tissue can herniate out of the skull through existence of a weak point between the frontal and ethmoidal bones during embryonic development (Dhirawani et al., 2014). In human, frontoethmoidal meningoencephalocele is commonly found with other congenital defects of facial malformation such as cleft lip and cleft palate (Ganapathy et al., 2014). However, it is rarely diagnosed in domestic animals. Other congenital defects of the ears and eyes such as microtia, anotia (Beraud, 2012; Luquetti et al., 2012), microphthalmos, and anophthalmos (Dell, 2010; Shibuya et al., 2000) are also rarely found in dogs, whereas congenital malformations of vertebrae and ribs are frequently found, especially in brachycephalic dogs such as pug and French bulldog (Gutierrez-Quintana et al., 2014). This case will describe the characteristics of frontoethmoidal meningoencephalocele with multiple congenital anomalies of facial and vertebral malformation in a stillborn puppy.

## Materials and Methods

**Specimen and anatomical study:** After full term of first pregnancy, a one-and-a-half-year-old Shih-Tzu bitch previously mated with the same breed went into natural labor at N.P. Animal Hospital, Bangkok, Thailand. One of four puppies had facial abnormalities and died at birth. It was preserved in 10% buffered formalin for further morphological studies. Body length and weight were measured and pictures were taken.

**Histopathological study:** Briefly, a membrane-bound facial mass located at the center of the puppy's face was removed, cut through a mid-sagittal plane, following the standard protocol for histopathological tissue sectioning, and stained with H&E.

**Radiographic study:** Total body radiographs were taken in lateral and dorsoventral positions using Universal® X-ray machine model MoD3486-50 at 75 kV, 1/50 second.


## Results and Discussion

The puppy was 12 centimeters long and weighed 104 grams. Gross anatomical study revealed anophthalmia of the left eye and microphthalmia of the right eye (Fig. 1a-c). Anophthalmia is characterized by the absence of the globes and ocular tissue within the orbits (Shibuya et al., 2000). The association of anophthalmos with other congenital defects including facial asymmetry and ear deformities is commonly found (Dell, 2010; Samson and Viljoen, 1995). During the first 3 weeks of gestation, optic vessels grow and stimulate neuro-ectodermal tissue to proliferate and form lens placode. Anophthalmos is caused by the stop of optic vessel growth, resulting in no formation of ocular structures (Dell, 2010). Microphthalmos is an eye with size reduction because the growth of optic cup is inhibited. Several factors might cause microphthalmia such as genetic disorder, uterine infection, drug toxicities, and maternal vitamin A deficiency (Kennelly et al., 2011; Tsutsui et al., 1993; Verma and Fitzpatrick, 2007; Williamson and FitzPatrick, 2014). Recently, bilateral microphthalmos has been reported in a Pomeranian puppy (Dell, 2010). Our Shih-tzu puppy had both microtia of the right ear, characterized by underdevelopment of pinna, and anotia of the left ear, characterized by completely undeveloped pinna, which might be associated with abnormalities of the external auditory canal and middle ear (Luquetti et al., 2012) (Fig. 1a-c). Microtia in human has been reported many times and this anomaly was demonstrated for the first time in a 2-month-old male mixed breed dog (Rezaei et al., 2015). In addition, anotia has been reported to be associated with distinct Mendelian autosomal recessive syndrome in St. Bernard (Villagomez and Alonso, 1998). This evidence supports that, at least, anotia is related to genetic disorder.


Interestingly, a mass with the size of 1.4 x 2 cm covered by a clear membrane was found at the center of the face, causing disappearance of the nose and appearance of large cleft lip (Fig. 1a-c). Radiographs revealed incomplete formation of maxillary and frontal bones where the mass protrusion occurred (Fig. 1d). The incomplete frontal bone formation is shown in Fig. 2a after the skin of the puppy's head was dissected. Nasal cavity, nasal septum, ethmoid bone and hard palate were not observed after removal of the entire brain with the mass as shown in Fig. 2b. Meningoencephalocele in our case involved intranasal cavity, similar to most cases of live dogs in a recent report (Lazzerini et al., 2017). H&E staining of mid-sagittal tissue section revealed that the mass was composed of numerous neurons, microglia, and capillaries surrounded by astrocytes. The membrane which encapsulated the mass consisted of fibroblasts and connective tissue (Fig. 3a-b). Encephalocele is characterized by a protrusion of cranial content throughout the cranium according to congenital defects of the skull. In case, herniation containing both brain matter and meninges is called meningoencephalocele. It has been identified into several classifications depending on the location of the skull defect. These include occipital, frontoethmoidal, basal, and cranial vault types, which were proposed by Matson (Fountas et al., 2005). In humans, patients with

frontoethmoidal meningoencephalocele had swelling over the bridge of nose or inner canthus of the eye since birth (Suwanwela and Suwanwela, 1972). Using these criteria, the stillbirth puppy in our study had frontoethmoidal meningoencephalocele, which is rarely found in dogs as opposed to Southeast Asian people, in which it is commonly found with an incidence of 1:5000 of new born babies (David, 1993; Dhirawani et al., 2014). According to history taking, the cause of the abnormality seems not to be from using drugs or being exposed to chemical agents of the bitch. Recently, ethmoidal encephalocele in a 6-month-old cross-bred dog and intranasal meningoencephalocele



in a 5-month-old female border collie have been reported. Both dogs showed sign of seizure from cerebral dysfunction (Jeffery, 2005; Martle et al., 2009). The etiology of frontoethmoidal meningoencephalocele is still unknown. However, failure of the neural plate to transform to the neural tube caused by incomplete separation between the surface ectoderm (epithelium) and neuroectoderm during early gestation period is a possible cause of the defect (Hoving, 2000). Besides genetics, environmental factors such as fungal infection and teratogenic agents can be the causes of encephalocele (Dhirawani et al., 2014).



**Figure 1** (a) The puppy's face (front view) showing frontoethmoidal meningoencephalocele (black arrow) and cleft lip (white arrow). (b) The puppy's face (left side) showing frontoethmoidal meningoencephalocele (\*) and cleft lip (black arrow) under the mass. (c) The puppy's face (right side) showing microphthalmos (white arrow), microtia (black arrow), and frontoethmoidal meningoencephalocele (\*). (d) Radiograph (lateral position) illustrating frontoethmoidal meningoencephalocele (white arrowhead) and incomplete formation of the frontal bone (white arrow). (e) Radiograph (dorsoventral position) showing incomplete formation of the frontal bone (white arrowhead), butterfly lumbar vertebrae (black arrowhead), rib fusion (black and white arrows). R = right side. (f) Radiograph (dorsoventral position) revealing lateral aplasia of the 2<sup>nd</sup> and 3<sup>rd</sup> cervical vertebrae (white arrow). White arrowhead pointing at frontoethmoidal meningoencephalocele. R = right side.



**Figure 2** (a) The puppy's head with opening skin showing frontal bone (F) with incomplete formation area (arrowhead). FEEM was still intact (Star). Scale bar = 1 cm. (b) By cutting through the frontal bone, the entire brain and FEEM were removed. Associated structures including nasal cavity, nasal septum, ethmoid bone and hard palate were not observed. T = tongue, F = frontal bone, M = maxilla bone covered with gum and teeth, B = part of brain. Scale bar = 1 cm.



**Figure 3** (a) Histopathological sections of the membrane-bound mass protruding through the face (frontoethmoidal meningoencephalocele) showing several parts of the brain, meninges, nasal, and palatine structures. (b) Higher magnification of (a) showing numerous neurons, microglia, astrocytes, and capillaries. Magnification (a) 20x; (b) 400x.

Additionally, lateral aplasia of the 2<sup>nd</sup> and 3<sup>rd</sup> cervical vertebrae and ventral and median aplasia of the 3<sup>rd</sup> to 13<sup>th</sup> thoracic or butterfly vertebrae were observed. The shape of both left and right ribs was abnormal with fusion on some parts of the right side. Incomplete development of the 7<sup>th</sup> lumbar vertebra was clearly seen (Fig. 1e-f). Possible mechanisms responsible for the malformation of vertebrae include absence of vertebral vascularization, teratogenic effects, and genetic abnormalities (Westworth and Sturges, 2010).

Taken together, the stillbirth puppy in this report had multiple congenital anomalies including frontoethmoidal meningoencephalocele, anotia, microtia, anophthalmia, microphthalmia, cleft lips, and vertebral malformations. It can be conclude that, at least, genetic disorders may be involved. The developmental defects of this puppy probably occurred during the first 3 weeks of gestation and they were caused by the defective process of neuroectodermal development.

### Acknowledgements

We would like to thank Dr. Pattana Rattanashinnakorn, the director of N.P. Animal Hospital for sample provision, and also Dr. Suwit Kumtorntip, Dr. Siripong Kiatkittikun, and radiology staff from Small Animal Hospital of Chulalongkorn University for their help in radiographic facilities. We are grateful to Asst. Prof. Dr. Nan Choisunirachorn for her advantage comments in radiographic interpretation, Assoc. Prof. Dr. Weerapong Koykul for his English correction, Mr. Pakorn Prajammuang, Assoc. Prof. Dr. Rabin Rattanapani, Assoc. Prof. Dr. Payattra and Assoc. Prof. Dr. Nongluck Tantilipikara for their valuable comments in histological discussion.

### References

Beraud R 2012. Surgical management of 2 different presentations of ear canal atresia in dogs. *Can Vet J.* 53(4): 412-418.

David DJ 1993. Cephaloceles: classification, pathology, and management--a review. *J Craniofac Surg* 4(4): 192-202.

Dell M 2010. Severe bilateral microphthalmos in a Pomeranian pup. *Can Vet J* 51(12): 1405-1407.

Dhirawani RB, Gupta R, Pathak S and Lalwani G 2014. Frontoethmoidal encephalocele: Case report and review on management. *Ann Maxillofac Surg* 4(2): 195-197.

Fountas KN, Smith JR, Jenkins PD and Murro AM 2005. Spontaneous motor cortex encephalocele presenting with simple partial seizures and progressive hemiparesis. Case report and review of the literature. *Neurosurg Focus* 19(3): E10.

Ganapathy A, T S, Swer MH and Rao S 2014. Occipital meningoencephalocele with Cleft Lip, Cleft Palate and Limb Abnormalities- A Case Report. *J Clin Diagn Res.* 8(12): AD03-05.

Gutierrez-Quintana R, Guevar J, Stalin C, Faller K, Yeamans C and Penderis J 2014. A proposed radiographic classification scheme for congenital thoracic vertebral malformations in brachycephalic "screw-tailed" dog breeds. *Vet Radiol Ultrasound.* 55(6): 585-591.

Holmes AD, Meara JG, Kolker AR, Rosenfeld JV and Klug GL 2001. Frontoethmoidal encephaloceles: reconstruction and refinements. *J Craniofac Surg.* 12(1): 6-18.

Hoving EW 2000. Nasal encephaloceles. *Childs Nerv Syst* 16(10-11): 702-706.

Jeffery N 2005. Ethmoidal encephalocoele associated with seizures in a puppy. *J Small Anim Pract* 46(2): 89-92.

Kennelly K, Brennan D, Chummun K and Giles S 2011. Histological characterisation of the ethanol-induced microphthalmia phenotype in a chick embryo model system. *Reprod Toxicol* 32(2): 227-234.

Lazzerini K, Gutierrez-Quintana R, Jose-Lopez R, McConnell F, Goncalves R, McMurrrough J, De Decker S, Muir C, Priestnall SL, Mari L, Stabile F, De Risio L, Loeffler C, Tauro A, Rusbridge C, Rodenas S, Anor S, de la Fuente C, Fischer A, Bruehschwein A, Penderis J and Guevar J 2017. Clinical Features, Imaging Characteristics, and Long-term Outcome of Dogs with Cranial Meningocele or Meningoencephalocele. *J Vet Intern Med* 31(2): 505-512.

Luquetti DV, Heike CL, Hing AV, Cunningham ML and Cox TC 2012. Microtia: epidemiology and genetics. *Am J Med Genet A* 158A(1): 124-139.

Martle VA, Caemaert J, Tshamala M, Van Soens I, Bhatti SF, Gielen I, Piron K, Chiers K, Tiemessen I and Van Ham LM 2009. Surgical treatment of a canine intranasal meningoencephalocele. *Vet Surg* 38(4): 515-519.

Rezaei M, Mahmoudi T, Ebrahimi M and Vosugh D 2015. First report of microtia in dog. *Comp Clin Pathol.* 24: 699-702.

Samson G and Viljoen D 1995. A case of lateral facial cleft, cleft lip and palate, anophthalmia, microtia, clavicular agenesis and asternia. *Clin Dysmorphol* 4(3): 251-254.

Shibuya K, Tajima M and Nunoya T 2000. Anophthalmia and retinal degeneration associated with stenosis of the optic foramen in Fischer 344 rats. *Vet Pathol* 37(3): 264-267.

Suwawela C and Suwanwela N 1972. A morphological classification of sincipital encephalomeningoceles. *J Neurosurg* 36(2): 201-211.

Tsutsui Y, Kashiwai A, Kawamura N and Kadota C 1993. Microphthalmia and cerebral atrophy induced in mouse embryos by infection with murine cytomegalovirus in midgestation. *Am J Pathol* 143(3): 804-813.

Verma AS and Fitzpatrick DR 2007. Anophthalmia and microphthalmia. *Orphanet J Rare Dis* 2: 47.

Villagomez DA and Alonso RA 1998. A distinct Mendelian autosomal recessive syndrome involving the association of anotia, palate agenesis, bifid tongue, and polydactyly in the dog. *Can Vet J* 39(10): 642-643.

Westworth DR and Sturges BK 2010. Congenital spinal malformations in small animals. *Vet Clin North Am Small Anim Pract* 40(5): 951-981.

Williamson KA and FitzPatrick DR 2014. The genetic architecture of microphthalmia, anophthalmia and coloboma. *Eur J Med Genet* 57(8): 369-380.

## บทคัดย่อ

### โรคง่วงซึ้งและความผิดปกติแต่กำเนิดของใบหน้าและกระดูกโครงร่างในลูกสุนัขสายแรกคลอด

พรหมพร รักษาเสรี<sup>1</sup> วุฒิชัย กลมเกลี้ยง<sup>1\*</sup>

แม่สุนัขพันธุ์ชิลล์สุอายุ 1 ปีครึ่ง ท้องแรก คลอดลูก 4 ตัวโดยวิธีธรรมชาติเมื่อครรภ์กำหนด มีลูกเพศผู้ 1 ตัว น้ำหนัก 104 กรัม ยาว 12 เซนติเมตร สายแรกคลอด พร้อมกับความผิดปกติหลายอย่างของใบหน้าและกระดูกโครงร่าง จึงเก็บรักษาในฟอร์มมาลิ่น 10% เพื่อทำการศึกษาทางมหาวิทยาลัยวิทยาและรังสีวิทยา และรังสีวิทยาต่อไป การศึกษาทางมหาวิทยาลัยวิทยาพบความผิดปกติแต่กำเนิดของตา โดยไม่มีตาชาญ ส่วนหัววัวมีขนาดเล็กกว่าปกติ พบรความผิดปกติของหู โดยไม่มีหูชาญ ส่วนหูวัวมีขนาดเล็กกว่าปกติ พบรปภะแห่งที่เด่นชัดคือพบก้อนเนื้อสีน้ำตาลเมื่อยื่นบางใส่หูมโน่อกมากกลางใบหน้าแทนที่จะมูก ซึ่งเมื่อนำไปศึกษาทางจุลพยาธิวิทยาพบว่าประกอบด้วยเนื้อเยื่ออของระบบประสาท ส่วนกล้ามได้แก่ เซลล์ประสาทและเซลล์ค้าจูนชนิดต่าง ๆ รวมถึง microglia และ astrocytes การถ่ายภาพรังสีทั้งทั่วพบการเจริญที่ไม่สมบูรณ์ ของกระดูก frontal bones ในตำแหน่งที่พับก้อนเนื้อโผล่มากกลางใบหน้า หัวหมดนี้ยื่นยันลักษณะของโรคง่วงซึ้ง (frontoethmoidal meningoencephalocele) นอกจากนี้ ยังพบกระดูกคลอกห่อนที่ 1-3 กระดูกคลอกห่อนที่ 3 ลงไป รวมทั้งกระดูกเอวทั้งหมดมีรูปร่างผิดปกติ กระดูกซี่โครงมีรูปร่างผิดปกติทั้งด้านซ้ายและขวา ในขณะที่มีการเชื่อมกันของกระดูกซี่โครงด้านขวาหลายซี่ การศึกษาครั้งนี้เป็นการแสดงภาวะง่วงซึ้งในลูกสุนัขสายแรกคลอดในประเทศไทยที่เกิดร่วมกับความผิดปกติหลายอย่างรวมทั้งใบหน้าและกระดูกโครงร่าง ซึ่งถือเป็นภาวะที่พบได้น้อยมาก

**คำสำคัญ:** โรคง่วงซึ้ง ใบหน้าผิดปกติ กระดูกโครงร่างร่วมผิดปกติ เป็นแต่กำเนิด สุนัข

<sup>1</sup>ภาควิชาการวิภาคศาสตร์ คณะสัตวแพทยศาสตร์ จุฬาลงกรณ์มหาวิทยาลัย ปทุมวัน กรุงเทพฯ 10330

\*ผู้รับผิดชอบบทความ E-mail: Wuthichai.K@chula.ac.th