

Monitoring Ovarian Cycles by Fecal Progesterone Analysis in Thamin Eld's Deer Hinds (*Rucervus eldii thamin*)

Orasa Phraluk^{1,2} Chainarong Punkong³ Ampika Thongphakdee⁴

Boripat Siriaroonrat⁴ Worawidh Wajjwalku⁵ Nikorn Thongtip^{1,2,6*}

Abstract

The aims of this study were to demonstrate the effectiveness of fecal progesterone metabolite monitoring technique during estrus synchronization and to investigate fecal progesterone metabolite profile in Eld's deer (*Rucervus eldii thamin*) hinds for one year (February 2009-January 2010). Fecal samples from five hinds were collected weekly, subjected to hormone extraction and subsequently progesterone analysis using enzyme immunoassay (EIA). Estrus synchronization and artificial insemination were performed during breeding season (June 2009). This included an intramuscular injection of ProstaglandinF_{2α} (5 mg) in combination with a vaginal insertion of a controlled internal drug-releasing device (CIDR type G) for 14 d. After CIDR-G removal for 70 h, all hinds were inseminated with frozen thawed semen by laparoscopic technique. This study found that one from five hinds responded to hormone stimulation. The fecal progesterone metabolite of the responded hind (no.11) was significantly highest compared with the non-responded hinds (no.7, no.42, no.43 and no.44) during the CIDR-G insertion (2,360±320 ng/g dry feces) ($p<0.05$), whereas the fecal progesterone concentrations of the four non-responded hinds during synchronization did not differ from basal progesterone concentrations ($p>0.05$). Mean±SE of progesterone metabolite concentrations during March to August (390.1±33.2 ng/g) were significantly higher than during September to February (291.4±20.1 ng/g) ($p<0.05$). The non-pregnant Eld's deer hinds exhibited multiple estrous cycles. Long estrous cycle (24.9±2.2 d) was observed throughout the year and mean number of estrous cycles was 14.0±0.4 cycles/animal/year. The concentration (mean±SE) of fecal progesterone during gestation was 816.9±60.0 ng/g in early pregnancy, 1,998.1±416.7 ng/g in middle pregnancy and 5,995.5±757.4 ng/g in late pregnancy. Consequently, fecal steroid analysis by EIA for monitoring of reproductive function and fecal steroid measurement can be used as an important breeding management in Eld's deer.

Keywords: Eld's deer, enzyme immunoassay, hormone, progesterone

¹Center for Agricultural Biotechnology, Kasetsart University, Kamphaeng Saen Campus, Nakhon Pathom 73140, Thailand

²Center of Excellence on Agricultural Biotechnology, Science and Technology Postgraduate Education and Research Department Commission on Higher Education, Ministry of Education (AG-BIO/PERDO-CHE) Bangkok 10900, Thailand

³Department of Conservation, Research and Animals Health, Khao Kheow Open Zoo, Zoological Park Organization, Chonburi 20110, Thailand

⁴Bureau of Conservation, Research and Education, Zoological Park Organization, Bangkok 10300, Thailand

⁵Department of Pathology, Faculty of Veterinary Medicine, Kamphaeng Saen Campus, Nakhon Pathom 73140, Thailand

⁶Department of Large Animal and Wildlife Clinical Sciences, Faculty of Veterinary Medicine, Kamphaeng Saen Campus, Nakhon Pathom 73140, Thailand

*Correspondence: foetnit@ku.ac.th

Introduction

Eld's deer (*Rucervus eldii*) is a subtropical cervid species of South and Southeast Asia. This deer species is listed as endangered by the IUCN (Timmins and Duckworth, 2008), and is one of fifteen species listed in the Thailand's National Wildlife Reservation and Protection Act since 1992. Two subspecies of Eld's deer; the Thamin Eld's deer (*Rucervus eldii thamin*) and Siamese Eld's deer (*Rucervus eldii siamensis*) (Balakrishnan et al., 2003), historically existed in Thailand's dry forest and were extinct in the wild since 1980s. However, both populations are maintained in captivity. Because of a small population size and low heterozygosity, Siamesis population exhibits low fecundity and high incidence of stillbirths (Personal communication, Dusit Zoo, Thailand). From the joint conservation effort among Thai government agencies, Zoological Park Organization, Department of National Parks, Wildlife and Plant Conservation, academics and Smithsonian Conservation Biology Institute, the Thamin Eld's deer have been successfully bred and reintroduced into the wild (Buranapim et al., 2008; Prempreet et al., 2013). At present, a total of 691 Eld's deer (641 Thamin and 50 Siamesis) are being maintained in *ex-situ* conservation centers (unpublished data, Nikorn Thongtip).

Female Eld's deer attain puberty at 2 years of age (Myint et al., 2001). Data from behavioral (Wemmer and Grodinsky, 1988) and endocrine studies (Monfort et al., 1990a; Monfort et al., 1990b) indicate that the females are seasonally polyestrous with the length of estrous cycle ranging from 13.8 to 21.5 days (Hosack et al., 1997; Monfort et al., 1990a). Even though the animals were translocated to Northern temperate latitudes, estrous behaviors occur in early spring (January-March) and anestrus begin in autumn (August-October) (Monfort et al., 1990b). Gestational period of Eld's deer is 235-245 d (Monfort et al., 1993). In captivity, calves usually born between September and November in France (48°N latitude, Prescott, 1987) and the United States (38°N latitude, Wemmer and Grodinsky, 1998) and September to February in China (19.6°N latitude, Song and Zeng, 2003).

It is well recognized that captive breeding programs play important roles in conserving wildlife. These populations are important as a resource for public awareness programs in zoos and as a hedge in case of some catastrophic events (e.g., disease) occurring in nature. In many wildlife species that maintained *ex situ*, assisted reproductive technology has been widely utilized to maximize individual's reproductive performance and to maintain genetic diversity within the population. Non-invasive endocrine monitoring is one of the useful tools for breeding management. It has been used to generate information on species, reproductive biology, knowledge that is essential for successful establishment of assisted reproductive technologies such as artificial insemination. There has been a report on the monitoring of the urinary steroid pregnanediol-3 α -glucuronide in captive of Eld's deer hinds. It indicates that Eld's deer has prolonged ovarian activity and exhibits serial reproductive cycle in one year at temperate latitudes. Currently, information about

reproductive cycle of Eld's deer in native habitat is limited. Non-invasive monitoring of reproductive status has been studied in several deer species including white-tailed deer (*Odocoileus virginianus*) (Kapke et al., 1999), brown brocket deer (*Mazama gouazoubira*) (Pereira et al., 2006), Pere David's deer (*Elaphurus davidianus*) (Li et al., 2001) and Sika deer (*Cervus nippon*) (Yamauchi et al., 1999). Two objectives of this study were to utilize fecal progesterone monitoring in examining the response to an estrus synchronization protocol and to assess seasonal fluctuation of fecal steroid concentrations in the Eld's deer hinds.

Materials and Methods

Animals and sample collection: Thamin Eld's deer hinds (n=5, 3-6 years old; 35-45 kg body weight) were maintained at Kasetsart University Kampheng Sean Campus, Thailand (14°N latitude). All hinds were housed in stocking rate and exposed to natural photoperiod. In this study, there was no stag as a teaser animal. They were fed *ad libitum* fresh Paragrass (*Brachiaria mutica*), commercial pellets, mineral block and water supply. Fecal samples were collected weekly during February 2009 to January 2010. The fecal samples were collected at 8:30 weekly. All samples were stored at -20°C until steroid analysis.

Synchronization of estrus and artificial insemination: All Eld's deer hinds were synchronized with treatment consisting of 5 mg PGF_{2 α} analogue (PGF_{2 α} 5 mg/ml/hind; Lutalyse™, Pfizer Animal Health, U.S.A.) on day 0 and a CIDR-G (Eazi-Breed™ Sheep Insert, Pfizer Animal Health, New Zealand) was inserted intravaginally into each restrained hind for 14 d. CIDR-G was pulled out on day 14. After CIDR-G removal for 70 h (day 17), each hind was artificially inseminated by using laparoscopic procedures described for Eld's deer (Monfort et al., 1993). Total of post-thaw sperm number for insemination was 25 x 10⁶ spermatozoa with >50% progressive motility.

Pregnancy diagnosis and fawning data: Pregnancy was diagnosed by progesterone measurement. Behavior and changing of physiology of all hinds were observed daily throughout the experiment.

Fecal steroid hormone extraction and enzyme immunoassay: Frozen fecal samples were thawed and dried at 60°C for 72 h. Dried fecal samples were ground. Steroid hormones were extracted with ethanol based procedure as described by Brown et al. (2004). Briefly, 0.1g feces was mixed with 5 ml of 90% ethanol in a 15 ml glass tube after vortexing. The tubes were then boiled in a water bath at 90°C for 20 min, and centrifuged at 1,500 rpm for 20 min. Supernatant was transferred into a new sample tube. A total of 5 ml of 90% ethanol was added to a pellet, vortexed for 30 seconds and recentrifuged for 15 min. The second supernatant was combined with the first one, dried and reconstituted in 1 ml of dilution buffer for frozen storage. Progesterone EIA was performed as described by Brown et al. (2004). Fecal samples were assayed in duplicate. Antibody for progesterone metabolite analysis (monoclonal pregnane CL no.425, 1:10,000

dilution) was obtained from Coralie Monro (University of California-Davis, CA, USA). Dosages of fecal extract samples from the Eld's deer were conducted according to Brown et al. (2004), by observation of parallelism between serial dilution (1:2-1:2,048) and standard curve (7.8-2,000 pg/ml). The samples were diluted to 1:200 for non-pregnant Eld's deer and 1:400 for pregnant Eld's deer in assay buffer. Sensitivity of assay was 7.8 pg/ml. Intra- and inter assay coefficients of variation were 6.77% and 6% (n=11 plates).

Statistical analysis: Data were presented as mean±SE. The effect of estrus synchronization with PGF_{2α} injection and CIDR-G insertion was evaluated by comparing fecal progesterone concentrations before, during CIDR-G insertion and after removal using paired samples T-test. The lowest progesterone concentrations of each cycle of each hind were defined as baseline progesterone and it was used for calculation. The baseline progesterone was calculated using an interactive process in which values that exceeded the mean plus 1.5 standard deviation were excluded. Average was then recalculated and the elimination process was repeated until no value exceeded the mean plus 1.5 standard deviation (Brown et al., 2001). If the values were more than the baseline progesterone value, they were considered as a luteal phase, whereas if the values were less than the baseline progesterone values, they were considered as an inter-luteal phase. The interval between progesterone concentration of luteal and inter-luteal phases was considered as duration of anestrus cycle (Pereira et al., 2006; Zanetti et al., 2010). Progesterone concentrations of luteal phase and inter-luteal phase were compared by paired samples T-test. Breeding season of Eld's deer in Thailand begins in March to August and non breeding season starts in September to February. Differences in fecal progesterone concentrations of the non-pregnant hinds during March-August and September-February were compared by using two samples T-test. Mean±SE of progesterone concentrations in early (1-11 weeks), middle (12-21 weeks) and late pregnancy (22-32 weeks) were determined by aligning the gestation day. Differences among the three gestation periods were determined using Kruskal-Wallis test, followed by Duncan's new multiple range test. Values of $p < 0.05$ were considered significant.

Results

Progesterone concentrations of estrous synchronized hinds are shown in Table 1. The data from five estrous synchronized hinds in this study revealed that four hinds (no.7, no.42, no.43 and no.44) did not respond to CIDR insertion since there were no variations in fecal progesterone metabolites. For the remaining hind (no.11), the progesterone concentrations increased during CIDR insertion, but then declined to similar level before hormone treatment.

Fecal progesterone concentration ranged from 50 to 1,640 ng/g in non-pregnant hinds. Progesterone metabolite concentrations profiles of individual Eld's deer hinds are shown in Fig 1. Mean±SE of progesterone during March to August (390.1±33.2 ng/g of dry feces) was significantly higher than that during September to February (291.4±20.1 ng/g of dry feces) ($p < 0.05$) (Fig 3). Mean estrous cycle duration was 24.9±2.2 d. Progesterone concentration of the luteal phase was higher ($p < 0.05$) than the inter-luteal phase. Mean of progesterone concentration in the inter-luteal phase and the luteal phase were 176.8±12.8 and 391.0±23.6 ng/g, respectively. Mean number of estrous cycles in a year was 14.0±0.4 cycles (range, 13-15). After artificial insemination, 1 from 5 hinds (no.11) was pregnant. The pregnant hind increased her feed intake and was isolated from the other hind. Before the fawn was to be born, the pregnant hind searched a safe place to give birth and hide her fawn from predators. Gestation duration was 242 d (34.5 weeks). The female fawn was born alive unassisted. Fecal progesterone level for the pregnant hind is presented in Fig 3. Difference in progesterone concentrations among early pregnancy, middle pregnancy and late pregnancy were significant ($p < 0.05$). Mean±SE of weekly progesterone profile in early pregnancy was 816.9±60.0 ng/g of dry feces, in middle pregnancy was 1,998.2±416.8 ng/g of dry feces and in late pregnancy was 5,995.6±757.4 ng/g of dry feces. Fecal progesterone metabolite levels were maintained between week 1 to 15 (average 659.3±78.0 ng/g of dry feces) before markedly increased during week 16-32 (4530.3±651.4 ng/g of dry feces). The highest peak was found on week 32 (12,160 ng/g of dry feces).

Table 1 Mean±SE fecal progesterone metabolite concentration before, during and after CIDR-G removal in each Eld's deer hind

| Events* | Mean ±SE of fecal progesterone metabolite concentration (ng/g of dry feces) | | | | |
|---|---|------------------------|---------|--------|--------|
| | No.7 | No.11 | No.42 | No.43 | No.44 |
| Before CIDR-G insertion (2 weeks: n = 2) | 770±330 ^b | 640±220 ^a | 390±250 | 310±30 | 140±60 |
| During CIDR-G insertion (2 weeks: n = 2) | 1,070±130 ^b | 2,360±320 ^b | 160±20 | 490±90 | 180±20 |
| CIDR-G removal (2 weeks: n = 2) | 270±70 ^a | 550±150 ^a | 260±20 | 200±40 | 120±20 |

^{a,b}Different subscripts within column are significantly different at the $p < 0.05$ level.




Figure 1 Individual progesterone metabolite concentrations of non-pregnant Eld's deer hinds (no.K7, K42, K43, K44) (Black bars = CIDR insertion, arrow = time of artificial insemination)




Figure 2 Mean fecal progesterone metabolite concentration throughout the year in the pregnant Eld's deer, no.K11 (Black bars = CIDR insertion, arrow = time of AI)

Discussion

In this present study, the evaluation of fecal progesterone metabolite concentrations during estrus synchronization and one year monitoring of fecal progesterone metabolite concentration in the non-pregnant and the pregnant Eld's deer hinds using enzyme immunoassay were done. The estrus synchronization using $\text{PGF}_{2\alpha}$ injection is associated with exogenous progesterone (CIDR devices) for CL regression. The CIDR maintain exogenous progesterone in the blood circulation and excretes into

feces. After the removal of devices, normal estrous cycle restarts. The CIDR devices has been successfully performed in follow deer (*Dama dama*) (8/10; 80%) (Asher et al., 1990; Morrow et al., 1995), Brown brocket deer (*Mazama gouazoubira*) (3/3; 100%) (Zanetti et al., 2010). The obtained data in this present study represented the values of fecal progesterone concentrations during estrus synchronization. In deer no.7, no.42 no.43 and no.44, the concentrations did not differ from the normal luteal phase concentrations. The reason for poor response inducing of estrous might be caused by the ovarian activity at the time of CIDR insertion. The response of ovary on CIDR depended on the stage of estrous cycle at CIDR insertion and duration of treatment (Macmillan and Peterson, 1993). Macmillan et al (1991) reported that CIDR insertion in cattle during metestrus and diestrus might compromise endogenous progesterone production by the newly corpus luteum. Therefore, progesterone absorption from CIDR device may be varied (Thompson and Monfort, 1999; Rathbone et al., 2002). Two studies reported that plasma progesterone concentration of red deer hinds (*Cervus elaphus*) receiving single CIDR-G/CIDR-S insertion within the first 6 d (2-3 ng/ml) (Asher et al., 1993) was higher than plasma progesterone concentration of single CIDR-G insertion for 14 d (1.0 ng/ml) (Jopson et al., 1990).

Our data indicated that the Eld's deer hinds had multiple estrous cycles throughout the year. The average fecal progesterone concentration during March to August was higher than during September to February. Moreover, the fecal progesterone concentration patterns of this study were similar to the




Figure 3 Average of fecal progesterone metabolite concentrations during February 2009 to January 2010 in 4 Eld's deer hinds (K7, K42, K43, K44)

urinary pregnanediol-3 α -glucuronide (PdG) patterns of a previous report in the northern temperate latitudes (38°N latitude) (Hosack et al., 1997). Therefore, due to the higher progesterone concentration, it might be assumed that in Thailand during March to August was the breeding season. This was in accordance with a previous study reporting that the oocyte quality from the Eld's deer hinds in the breeding season had higher quality than during the non-breeding season in terms of number of follicles, follicle size, number of oocytes recovered and oocyte quality (Siriaroonrat, 2006).

The data of fecal progesterone profiles in this study coincided with the expression of endogenous rhythms in the Eld's deer reproductive hormones in northern temperate latitudes. It was not related to historical pattern of habitat resources, local food and photoperiod (Hosack et al., 1997). In this study, the mean of continuous estrous cycle (14.0 \pm 0.4 cycles) throughout 12 months was similar to a previous report in North America (13.8 \pm 0.5 cycles) (Hosack et al., 1997). However, in this present study, they had longer estrous cycles (24 d) than previous data (17.4-21.5 d for *thamin*; Monfort et al., 1990^a; 1990^b and 16.3 d for *hainanus*; Song and Zeng, 2003). However, these results were similar to the brown brocket deer (26.9 d) (Krepschi et al., 2013). The reason of longer estrous in this study remains unclear, probably the effect of stag absence on ovarian irregularity (Pereira et al., 2006). Therefore, we assumed that the longer estrous cycle of Eld's deer in this study might be related to the stag absence. However, further studies need to be done. The changes of fecal progesterone metabolite during pregnancy of the Eld's deer in this study were similar to previous data of urinary PdG concentrations pattern in pregnant Eld's deer hinds (Monfort et al., 1990^b; Monfort et al., 1993) and other deer species (Elk deer; White et al., 1995, Red brocket deer; Krepschi et al., 2013, Reindeer; Bubenik et al., 1997, Sika deer; Matsura et al., 2004, White tailed deer; Kapkeet al., 1999). The fecal progesterone level of pregnant hind was markedly higher than hinds that failed to conceive. The progesterone level slightly elevated in the initial stage of gestation because the *corpus luteum* is the only source of progesterone released in blood circulation for maintenance of pregnancy. In mid and late pregnancy, progesterone level is clearly much increased due to the

supplement source of progesterone by placenta (Flood et al., 2005).

In summary, fecal hormone analysis by using enzyme immunoassay could be useful for monitoring reproductive status in the Eld's deer such as estrous cycle, baseline hormone concentration, estrous synchronization and pregnancy diagnosis. Hormone concentration of individual female Eld's deer can be important information for Eld's deer conservation program.

Acknowledgements

This study was supported by the Center for Agricultural Biotechnology and the Center of Excellence on Agricultural Biotechnology (AG-BIO/PERDO-CHE), Kasetsart University. We would like to thank the staffs and organizations at Kasetsart University and the scientists and veterinarians of zoos under the Zoological Park Organization (Khao Kheow Open Zoo) for their support in this study. Moreover, we would like to thank Dr. Nucharin Songsasen from Smithsonian Conservation Biology Institute, USA for critical editing this manuscript.

References

- Asher GW, Fisher MW, Smith JF, Jabbour HN and Marrow CJ 1990. Temporal relationships between the onset, the preovulation LH surge and ovulation in farmed fallow deer, *Dama dama*. J Reprod Fert. 89:761-767.
- Asher GW, Fisher MW, Fennessy PF, Mackintosh CG, Jabbour HN and Marrow CJ 1993. Oestrous synchronization, semen collection and artificial insemination of farmed red deer (*Cervus elaphus*) and fallow deer (*Dama dama*). Anim Reprod Sci. 33:241-265.
- Balakrihnan CN, Monfort SL, Gaur A, Singh L and Sorenson MD 2003. Phylogeography and conservation genetics of Eld's deer (*Cervus eldi*). Mol Ecol. 12(1):1-10.
- Brown JL, Bellem AC, Michael F, Wildt DE and Roth TL. 2001. Comparative analysis of gonadal and adrenal activity in the Black and White rhinoceros in north America by noninvasive endocrine monitoring. Zoo Biology. 20: 483-486.
- Brown JL, Walker S and Steinman K 2004. Endocrine Manual for the Reproductive Assessment of Domestic and non-domestic species. 2nded. Endocrine Research Laboratory, Department of Reproductive Sciences, Conservation and Research Center, Nation Zoological Park Smithsonian Institution, Virginia, USA. 94 pp.
- Bubenik A, Schams D, White RJ, Rowell J, Blake J and Bartos L 1997. Seasonal levels of reproductive hormones and their relationship to the antler cycle of male and female reindeer (*Rangifer tarandus*). Comp Biochem Physiol. 116(2): 269-277.
- Buranapim N, Sukmasuang R and Bhumparkpan N 2008. Population Viability Analysis of Reintroducing Brown-antlered Deer (*Cervus eldi thamin*) in Huai Kha Khaeng Wildlife Sanctuary,

- Uthai Thane Province. Thesis. Graduate school. Kasetsart University.
- Flood PF, Tyler NJC, Read EK, Rodway MJ and Chedrese PJ 2005. Ovarian and placental production of progesterone and oestradiol during pregnancy in rein deer. *Anim Reprod Sci.* 85: 147-162.
- Hosack DA, Miller HV, Marchinton RL and Monfort SL 1997. Ovarian activity in captive Eld's deer (*Cervus eldi thamin*). *J Mammal.* 78(2):669-674.
- Jopson NB, Fisher MW and Suttie JM 1990. Endogenous and exogenous progesterone in reddeer hinds. *Anim Reprod Sci.* 23:61-73.
- Kapke CA, Arcese P, Ziegler TE and Scheffler GR 1999. Estradiol and progesterone metabolite concentration in white-tailed deer (*Odocoileus virginianus*) feces. *J Zoo Wildl Med.* 30:361-371.
- Krepschi VG, Polegato BF, Zanetti ES and Duarte JMB 2013. Fecal progestins during pregnancy and postpartum periods of captive red brocket deer (*Mazama americana*). [Online]. Available: <http://dx.doi.org/10.1016/j.anireprosci.2012.11.016>. Accessed June 20, 2014.
- Li C, Jiang Z, Jiang G and Fang J 2001. Seasonal changes of reproductive behavior and fecal steroid concentrations in Pere David's deer. *Hormones and Behavior.* 40:518-525.
- Macmillan KL, Taufu VK, Barnes DR and Day AM 1991. Plasma progesterone concentrations in heifers and cows treated with a new intravaginal device. *Anim Reprod Sci.* 26:25-40.
- Macmillan KL and Peterson AJ 1993. A new intravaginal progesterone releasing device for cattle (CIDR-B) for oestrous synchronization, increasing pregnancy rates and the treatment of post-partum anoestrous. *Anim Reprod Sci.* 33:1-25.
- Matsura Y, Sasamoto Y, Sato K, Takahashi Y, Suzuki M and Ohtaishi N 2004. Monitoring ovarian cycle and conception by fecal progesterone analysis in sika deer. *Ecol Res.* 19:397-404.
- Monfort SL, Wemmer C, Brown JL and Wildt DE 1990^a. Use of urinary hormone assays for evaluating endocrine patterns associated with the long-day breeding season in Eld's deer (*Cervus eldi*). *J Experimental Zoology.* 4:215-218.
- Monfort SL, Wemmer C, Kepler TH, Bush M, Brown JL and Wildt DE 1990^b. Monitoring ovarian function and pregnancy in Eld's deer (*Cervus eldi thamin*) by evaluating urinary steroid metabolite excretion. *J Reprod Fert.* 88:271-281.
- Monfort SL, Asher GW, Wildt DE, Wood TC, Schiewe MC, Williamson LR, Bush M and Rall WF 1993. Successful intrauterine insemination of Eld's deer (*Cervus eldi thamin*) with frozen thawed spermatozoa. *J Reprod Fert.* 99:459-465.
- Morrow CJ, Asher GW and Macmillan KL 1995. Oestrous synchronization in farmed Fallow deer (*Dama dama*): effect of season, treatment duration and the male on the efficacy of the intravaginal CIDR device. *Anim Reprod Sci.* 37:159-174.
- Myint A, Mashea WJ, Htung S, Tham A, Soe TM, Monfort S and Wemmer C 2001. Ecology and social organization of a tropical deer (*Cervus eldi thamin*). *J Mammal.* 83(3):837-471.
- Pereira RJG, Polegato BF, Souza S, Negrao JA and Duarte JMB 2006. Monitoring ovarian cycles and pregnancy in brown brocket deer (*Mazama gouazoubira*) by measurement of fecal progesterone metabolites. *Theriogenology.* 65:387-399.
- Prempre P, Sukmasuang R, Bhumparkphan N and Yindee M 2013. Post-released Monitoring of Hog deer, Eld's deer and Sambar deer in Salakphra Wildlife Sanctuary, Kanchanaburi Province.
- Rathbone MJ, Bunt CR, Ogle CR, Burggraaf S, Macmillan KL, Burke CR and Pikerling KL 2002. Re engineering of a commercially available bovine Intravaginal insert (CIDR insert) containing progesterone. *J Controlled Release.* 85:105-115.
- Siriaronrat B 2006. Development of *in vitro* fertilization for the endangered Eld's deer (*Cervus eldi thamin*): Effect of *in vitro* maturation, fertilization and culture on developmental competence. Thesis. (Philosophy), George Mason University, USA. 143 pp.
- Song YL and Zeng ZG 2003. Reproductive characteristics of female Eld's deer in their natural range in Hainan island, China. *Mammalia.* 67: 47-53.
- Thompson KV and Monfort SL 1999. Synchronization of oestrous cycles in sable antelope. *Anim Reprod Sci.* 57:185-197.
- Timmins RJ and Duckworth JW 2008. *Rucervus eldii* In: IUCN 2012: IUCN Red List of Threatened Species. Version 2012.1. [Online]. Available: iucnredlist.org. Accessed May 20, 2013.
- Wemmer C and Grodinsky C 1988. Reproduction in captive female brow-antlered Deer (*Cervus eldi thamin*). *J Mammal.* 69(2): 389-393.
- White PJ, Garrott RA, Kirkpatrick JF and Berkeley EV 1995. Diagnosing pregnancy in free-ranging elk using fecal steroid metabolites. *J Wildl Dis.* 31(4):514-522.
- Yamauchi K, Hamasaki S, Takeuchi Y and Mori Y 1999. Application of enzyme immunoassay to fecal steroid analysis in sika deer (*Cervus nippon*). *J Reprod Dev.* 45:429-434.
- Zanetti ES, Polegato BF and Duarte JMB 2010. Comparison of two methods of synchronization of estrus in brown brocket deer (*Mazama gouazoubira*). *Anim Reprod Sci.* 117: 266-274

บทคัดย่อ

การตรวจวงรอบการเป็นสัดด้วยการวิเคราะห์ปริมาณฮอร์โมนโปรเจสเตอโรนจากอุจจาระ ของละมั่งเพศเมียสายพันธุ์พม่า (*Rucervus eldii thamin*)

อรสา พระลักษณ์^{1,2} ชัยณรงค์ บั่นคง³ อัมพิกา ทองภักดี⁴ ปริพัตร ศิริอรุณรัตน์⁴

วรวิทย์ วัชชวัลคุ⁵ นิกร ทองทิพย์^{1,2,6*}

การศึกษานี้มีวัตถุประสงค์เพื่อตรวจหาระดับฮอร์โมนโปรเจสเตอโรนจากอุจจาระในช่วงเหนี่ยวนำการเป็นสัดเพื่อการทดลองผสมเทียมและศึกษาการเปลี่ยนแปลงของปริมาณฮอร์โมนโปรเจสเตอโรนที่พบในอุจจาระของละมั่งพันธุ์พม่าเพศเมีย (*Rucervus eldii thamin*) จำนวน 5 ตัวเป็นระยะเวลา 1 ปี (ก.พ. 2552-ม.ค. 2553) โดยเก็บตัวอย่างอุจจาระละมั่ง สัปดาห์ละ 1 ครั้ง จนกระทั่งเสร็จสิ้นการทดลองแล้วนำมาวิเคราะห์ปริมาณฮอร์โมนโปรเจสเตอโรนด้วยวิธี Enzyme immunoassay (EIA) ขั้นตอนการเหนี่ยวนำการเป็นสัดและการผสมเทียมดำเนินการในเดือนมิถุนายน ละมั่งทั้งหมดได้รับการฉีดฮอร์โมน PGF_{2α} (5 มก./ตัว) พร้อมกับสอดแท่งโปรเจสเตอโรนสังเคราะห์ (CIDR-G) ไว้ในช่องคลอดเป็นเวลา 14 วัน จึงถอดแท่งโปรเจสเตอโรนสังเคราะห์ออก ทำการผสมเทียมด้วยน้ำเชื้อแช่แข็งผ่านกลัองลาปาโรสโคป หลังจากถอดแท่งโปรเจสเตอโรนออกเป็นระยะเวลา 70 ชั่วโมง พบว่าปริมาณโปรเจสเตอโรนที่วิเคราะห์ได้ขณะที่มีการเหนี่ยวนำการเป็นสัดในละมั่งหมายเลข 11 มีค่าความเข้มข้นของฮอร์โมนโปรเจสเตอโรน (2,360±320 นาโนกรัม/กรัม) สูงกว่าละมั่งตัวอื่น (หมายเลข 7, 42, 43 และ 44) อย่างมีนัยสำคัญทางสถิติ ($p < 0.05$) ขณะเดียวกันไม่พบความแตกต่างของความเข้มข้นของฮอร์โมนโปรเจสเตอโรนในละมั่งทั้ง 4 ตัวระหว่างที่มีการเหนี่ยวนำการเป็นสัด ($p > 0.05$) ละมั่งสามารถแสดงวงรอบการเป็นสัดซ้ำหลายวงรอบ (multiple estrous cycle) ต่อปี มีความยาวของวงรอบการเป็นสัด 24.9±2.2 วัน คิดเป็น 14.0±0.4 รอบต่อปี ซึ่งค่าเฉลี่ยของฮอร์โมนโปรเจสเตอโรนในช่วงเดือนมีนาคมถึงเดือนสิงหาคม (390.1±33.2 นาโนกรัม/กรัม) มีค่าสูงกว่าช่วงเดือนกันยายนถึงเดือนกุมภาพันธ์ (291.4±20.1 นาโนกรัม/กรัม) อย่างมีนัยสำคัญทางสถิติ ($p < 0.05$) ส่วนละมั่งที่ตั้งท้องมีค่าเฉลี่ยของฮอร์โมนโปรเจสเตอโรนแบ่งออกเป็น 3 ระยะ ได้แก่ ระยะต้น (816.9±60.0 นาโนกรัม/กรัม), ระยะกลาง (1,998.1±416.8 นาโนกรัม/กรัม) และระยะท้ายของการอุ้มท้อง (5,995.6±757.4 นาโนกรัม/กรัม) ดังนั้นการใช้วิธีวิเคราะห์ปริมาณโปรเจสเตอโรนด้วยเทคนิค EIA สามารถประเมินการทำงานสรีรวิทยาของระบบสืบพันธุ์และเป็นเครื่องมือที่สำคัญในการจัดการด้านการสืบพันธุ์ละมั่งได้

คำสำคัญ: ฮอร์โมน โปรเจสเตอโรน ละมั่ง enzyme immunoassay

¹ศูนย์เทคโนโลยีชีวภาพเกษตร มหาวิทยาลัยเกษตรศาสตร์ วิทยาเขตกำแพงแสน จ.นครปฐม 73140 ประเทศไทย

²ศูนย์ความเป็นเลิศด้านเทคโนโลยีชีวภาพเกษตร สำนักพัฒนาบัณฑิตศึกษาและวิจัยด้านวิทยาศาสตร์และเทคโนโลยี สำนักงานคณะกรรมการอุดมศึกษา กรุงเทพฯ 10900

³ฝ่ายอนุรักษ์ วิจัย และสุขภาพสัตว์ สวนสัตว์เปิดเขาเขียว องค์การสวนสัตว์ในพระบรมราชูปถัมภ์ จ.ชลบุรี 20110 ประเทศไทย

⁴สำนักอนุรักษ์ วิจัยและการศึกษา องค์การสวนสัตว์ในพระบรมราชูปถัมภ์ กรุงเทพฯ 10300

⁵ภาควิชาพยาธิวิทยา คณะสัตวแพทยศาสตร์ มหาวิทยาลัยเกษตรศาสตร์ วิทยาเขตกำแพงแสน จ.นครปฐม 73140 ประเทศไทย

⁶ภาควิชาเวชศาสตร์คลินิกสัตว์ใหญ่และสัตว์ป่า คณะสัตวแพทยศาสตร์ มหาวิทยาลัยเกษตรศาสตร์ วิทยาเขตกำแพงแสน จ.นครปฐม 73140 ประเทศไทย

*ผู้รับผิดชอบบทความ E-mail: fvetnit@ku.ac.th